

The subdivision of this section is:

- References

References

Links are provided for those new references added to this edition (author name is highlighted in blue).

- Aan Het Rot M, Zarate Jr CA, Charney DS, Mathew SJ. Ketamine for depression: where do we go from here? *Biol Psychiatry* 2012;72:537-47.
- Adams JR, Drake RE, Wolford GL. Shared decision-making preferences of people with severe mental illness. *Psych Serv* 2007;58:1219-21.
- Alexopoulos GS, Abrams RC, Young RC, Shamoian CA. Cornell scale for depression in dementia. *Biol Psychiatry* 1988;23:271-84.
- Allan CL, Herrmann LL, Ebmeier KP. Transcranial magnetic stimulation in the management of mood disorders. *Neuropsychobiology* 2011;64:163-69.
- Alpert M, Silva RR, Pouget ER. Prediction of treatment response in geriatric depression from baseline folate level: interaction with an SSRI or a tricyclic antidepressant. *J Clin Psychopharmacol* 2003;23:309-13.
- American Psychiatric Association. *In Diagnostic and Statistical Manual of Mental Disorders DSM-5*. Fifth Edition. Washington, DC/London, England. 2013.
- American Psychiatric Association. *In Practice Guideline for the Treatment of Patients with Panic Disorder*. 2nd Edition, 2010.
- Anderson EL, Reti IM. ECT in pregnancy: a review of the literature from 1941 to 2007. *Psychosomatic Med* 2009;71:235-42.
- Anderson L, Lewis G, Araya R, et al. Self-help books for depression: how can practitioners and patients make the right choice? *Br J Gen Pract* 2005;55:387-92.
- Appelbaum PS. Assessment of patients' competence to consent to treatment. *N Engl J Med* 2007;357:1834-40.
- Appleton KM, Rogers PJ, Ness AR. Updated systematic review and meta-analysis of the effects of n-3 long-chain polyunsaturated fatty acids on depressed mood. *Am J Clin Nutr* 2010;91:757-70.
- Archer J, Bower P, Gilbody S, et al. Collaborative care for depression and anxiety problems. *Cochrane Database of Systematic Reviews* 2012(10):CD006525.
- Areán PA, Ayalon L, Hunkeler E, et al. Improving depression care for older, minority patients in primary care. *Med Care* 2005;43:381-90.
- Austin MP. To treat or not to treat: maternal depression, SSRI use in pregnancy and adverse neonatal effects. *Psychol Med* 2006;36:1663-70.
- Babor TF, Higgins-Biddle JC, Saunders JB, Monteiro MG. *Audit. In The Alcohol Use Disorders Identification Test: Guidelines for Use in Primary Care*. Second Edition. World Health Organization. 2001.
- Babyak M, Blumenthal JA, Herman S, et al. Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months. *Psychosom Med* 2000;62:633-38.
- Baik S, Gonzales JJ, Bowers BJ, et al. Reinvention of depression instruments by primary care clinicians. *Ann Fam Med* 2010;8:224-30.
- Bajbouj M, Merkl A, Schlaepfer TE, et al. Two-year outcome of vagus nerve stimulation in treatment-resistant depression. *J Clin Psychopharmacol* 2010;30:273-81.
- Ballenger JC, Davidson JRT, Lecrubier Y, et al. Consensus statement on depression, anxiety, and cardiovascular disease. *J Clin Psychiatry* 2001;62:24-27.

References

- Barca ML, Engedal K, Selbaek. A reliability and validity study of the Cornell scale among elderly inpatients, using various clinical criteria. *Dement Geriatr Cogn Disord* 2010;29:438-47.
- Baumann P, Nil R, Souche A, et al. A double-blind, placebo-controlled study of citalopram with and without lithium in the treatment of therapy-resistant depressive patients: a clinical, pharmacokinetic, and pharmacogenetic investigation. *J Clin Psychopharmacol* 1996;16:307-14.
- Beck A, Crain AL, Solberg LI, et al. Severity of depression and magnitude of productivity loss. *Ann Fam Med* 2011;9:305-11.
- Belnap BH, Kuebler J, Upshur C, et al. Challenges in implementing depression care management in the primary care setting. *Adm Policy Ment Health* 2006;33:65-75.
- Benedetti F, Colombo C, Pontiggia A, et al. Morning light treatment hastens the antidepressant effect of citalopram: a placebo-controlled trial. *J Clin Psychiatry* 2003;64:648-53.
- Bérard A, Ramos É, Rey É, et al. First trimester exposure to paroxetine and risk of cardiac malformations in infants: the importance of dosage. *Birth Defects Res B Dev Reprod Toxicol* 2007;80:18-27.
- Berlim MT, van den Eynde F, Tovar-Perdomo S, Daskalakis ZJ. Response, remission and drop-out rates following high-frequency repetitive transcranial magnetic stimulation (rTMS) for treating major depression: a systematic review and meta-analysis of randomized, double-blind and sham-controlled trials. *Psychol Med* 2014;44:225-39.
- Berman RM, Marcus RN, Swanink R, et al. The efficacy and safety of aripiprazole as adjunctive therapy in major depressive disorder: a multicenter, randomized, double-blind, placebo-controlled study. *J Clin Psychiatry* 2007;68:843-53.
- Biesheuvel-Leliefeld KE, Kok GD, Bockting CL, et al. Effectiveness of psychological interventions in preventing recurrence of depressive disorder: meta-analysis and meta-regression. *J Affect Disord* 2015;174:400-10.
- Birrer RB, Vemuri SP. Depression in later life: a diagnostic and therapeutic challenge. *Am Fam Physician* 2004;69:2375-82.
- Bloch MH, Hannestad J. Omega-3 fatty acids for the treatment of depression: systematic review and meta-analysis. *Mol Psychiatry* 2012;17:1272-82.
- Blomstedt P, Sjöberg RL, Hansson M, et al. Deep brain stimulation in the treatment of depression. *Acta Psychiatr Scand* 2011;123:4-11.
- Blumenthal JA, Lett HS, Babyak MA, et al. Depression as a risk factor for mortality after coronary artery bypass surgery. *Lancet* 2003;362:604-09.
- Bodkin JA, Lasser RA, Wines JD Jr., et al. Combining serotonin reuptake inhibitors and bupropion in partial responders to antidepressant monotherapy. *J Clin Psychiatry* 1997;58:137-45.
- Borges S, Chen YF, Laughren TP, et al. Review of maintenance trials for major depressive disorder: a 25-year perspective from the U.S. food and drug administration. *J Clin Psychiatry* 2014;75:205-14.
- Bortolotti B, Menchetti M, Bellini F, et al. Psychological interventions for major depression in primary care: a meta-analytic review of randomized controlled trials. *Gen Hosp Psychiatry* 2008;30:293-302.
- Bostwick JM, Pankratz VS. Affective disorders and suicide risk: a reexamination. *Am J Psychiatry* 2000;157:1925-32.
- Boukhris T, Sheehy O, Mottron L, Bérard A. Antidepressant use during pregnancy and the risk of autism spectrum disorder in children. *JAMA Pediatr* 2016;170:117-24.

References

- Bouwer C, Stein DJ. Buspirone is an effective augmenting agent of serotonin selective re-uptake inhibitors in severe treatment-refractory depression. *S Afr Med J* 1997;87:534-37.
- Bower P, Kontopantelis E, Sutton A, et al. Influence of initial severity of depression on effectiveness of low intensity interventions: meta-analysis of individual patient data. *BMJ* 2013;346: f540.
- Briggs JH, Bauer MS, McBride L, et al. Screening for thyroid disease in ambulatory patients with depression. *American Psychiatric Association Abstracts* NR144, 1993.
- Brown RL, Rounds LA. Conjoint screening questionnaires for alcohol and other drug abuse: criterion validity in a primary care practice. *Wis Med J* 1995;94:135-40.
- Brown C, Schulberg HC, Prigerson HG. Factors associated with symptomatic improvement and recovery from major depression in primary care patients. *Gen Hosp Psychiatry* 2000;22:242-50.
- Brown WA, Harrison W. Are patients who are intolerant to one serotonin selective reuptake inhibitor intolerant to another? *J Clin Psychiatry* 1995;56:30-34.
- Browne G, Steiner M, Roberts J, et al. Sertraline and/or interpersonal psychotherapy for patients with dysthymic disorder in primary care: 6-month comparison with longitudinal 2-year follow-up of effectiveness and costs. *J Affect Disord* 2002;68:317-30.
- Bruce MT, Ten Have TR, Reynolds III CF, et al. Reducing suicidal ideation and depressive symptoms in depressed older primary care patients: a randomized controlled trial. *JAMA* 2004;291:1081-91.
- Bull SA, Hu XH, Hunkeler EM, et al. Discontinuation of use and switching of antidepressants: influence of patient-physician communication. *JAMA* 2002;288:1403-09.
- Bush B, Shaw S, Cleary P, et al. Screening for alcohol abuse using the CAGE questionnaire. *Am J Med* 1987;82:231-35.
- Candy M, Jones L, Williams R, et al. Psychostimulants for depression (review). *Cochrane Database of Systematic Reviews* 2008(2):CD006722.
- Cappiello A, McDougle CJ, Malison RT, et al. Yohimbine augmentation of fluvoxamine in refractory depression: a single-blind study. *Biol Psychiatry* 1995;38:765-67.
- Chambers CD, Hernandez-Diaz S, Van Marter LJ, et al. Selective serotonin-reuptake inhibitors and risk of persistent pulmonary hypertension of the newborn. *N Engl J Med* 2006;354:579-87.
- Chen YM, Huang XM, Thompson R, Zhao YB. Clinical features and efficacy of escitalopram treatment for geriatric depression. *J Int Med Res* 2011;39:1946-53.
- Churchill R, Moore THM, Furukawa TA, et al. 'Third wave' cognitive and behavioural therapies versus treatments as usual for depression. *Cochrane Database Syst Rev* 2013(10)CD008705.
- Ciechanowski PS, Katon WJ, Russo JE. Depression and diabetes: impact of depressive symptoms on adherence, function, and costs. *Arch Intern Med* 2000;160:3278-85.
- Claassen CA, Trivedi MH, Rush AJ, et al. Clinical differences among depressed patients with and without a history of suicide attempts: findings from the STAR*D trial. *J Affect Disord* 2007;97:77-84.
- Coelho HF, Murray L, Royal-Lawson M, Cooper PJ. Antenatal anxiety disorder as a predictor of post-natal depression: a longitudinal study. *J Affect Disord* 2011;129:348-53.
- Cohen LS, Altshuler LL, Harlow BL, et al. Relapse of major depression during pregnancy in women who maintain or discontinue antidepressant treatment. *JAMA* 2006;295:499-507.
- Colangelo LA, He K, Whooley MA, et al. Higher dietary intake of long-chain w-3 polyunsaturated fatty acids is inversely associated with depressive symptoms in women. *Nutrition* 2009;25:1011-19.

References

- Cooney GM, Dwan K, Greig CA, et al. Exercise for depression. *Cochrane Database of Systematic Reviews* 2013(9)CD004366.
- Cooper C, Katona C, Lyketsos K, et al. A systematic review of treatments for refractory depression in older people. *Am J Psychiatry* 2011;168:681-88.
- Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression: development of the 10-item Edinburgh postnatal depression scale. *Br J Psychiatry* 1987;150:782-86.
- Cristancho P, Cristancho MA, Baltuch GH, et al. Effectiveness and safety of vagus nerve stimulation for severe treatment-resistant major depression in clinical practice after FDA approval: outcomes at 1 year. *J Clin Psychiatry* 2011;72:1376-82.
- Cuijpers P, Andersson G, Donker T, van Straten A. Psychological treatment of depression: results of a series of meta-analyses. *Nord J Psychiatry* 2011a;65:354-64.
- Cuijpers P, Dekker J, Hollon SD, Andersson G. Adding psychotherapy to pharmacotherapy in the treatment of depressive disorders in adults: a meta-analysis. *J Clin Psychiatry* 2009a;70:1219-29.
- Cuijpers P, Geraedts AS, van Oppen P, et al. Interpersonal psychotherapy for depression: a meta-analysis. *Am J Psychiatry* 2011b;168:581-92.
- Cuijpers P, Hollon SD, van Straten A, et al. Does cognitive behaviour therapy have an enduring effect that is superior to keeping patients on continuation pharmacotherapy? A meta-analysis. *BMJ Open* 2013;3:e002542.
- Cuijpers P, Karyotaki E, Andersson G, et al. The effects of blinding on the outcomes of psychotherapy and pharmacotherapy for adult depression: a meta-analysis. *Eur Psychiatry* 2015;30:685-93.
- Cuijpers P, Karyotaki E, Weitz E, et al. The effects of psychotherapies for major depression in adults on remission, recovery and improvement: a meta-analysis. *J Affect Disord* 2014a;159:118-26.
- Cuijpers P, Reynolds CF 3rd, Donker T, et al. Personalized treatment of adult depression: medication, psychotherapy, or both? A systematic review. *Depress Anxiety* 2012;29:855-64.
- Cuijpers P, Sijbrandij M, Koole SL, et al. Adding psychotherapy to antidepressant medication in depression and anxiety disorders: a meta-analysis. *World Psychiatry* 2014b;13:56-67.
- Cuijpers P, van Straten A, Andersson G, van Oppen P. Psychotherapy for depression in adults: a meta-analysis of comparative outcome studies. *J Consult Clin Psychol* 2008;76:909-22.
- Cuijpers P, van Straten A, Bohlmeijer E, et al. The effects of psychotherapy for adult depression are overestimated: a meta-analysis of study quality and effect size. *Psychol Med* 2010a;40:211-23.
- Cuijpers P, van Straten A, Schuurmans J, et al. Psychotherapy for chronic major depression and dysthymia: a meta-analysis. *Clin Psychol Rev* 2010b;30:51-62.
- Cuijpers P, van Straten A, Smit F, Andersson G. Is psychotherapy for depression equally effective in younger and older adults? A meta-regression analysis. *Int Psychogeriatr* 2009b;21:16-24.
- Cuijpers P, van Straten A, Warmerdam L. Behavioral activation treatments of depression: a meta-analysis. *Clin Psychol Rev* 2007;27:318-26.
- Cuijpers P, van Straten A, Warmerdam L, Andersson G. Psychotherapy versus the combination of psychotherapy and pharmacotherapy in the treatment of depression: a meta-analysis. *Depress Anxiety* 2009c;26:279-88.
- Dam J, Ryde L, Svejsø J, et al. Morning fluoxetine plus evening mianserin versus morning fluoxetine plus evening placebo in the acute treatment of major depression. *Pharmacopsychiatry* 1998;31:48-54.

References

- Danielsson L, Noras AM, Waern M, Carlsson J. Exercise in the treatment of major depression: a systematic review grading the quality of evidence. *Physiother Theory Pract* 2013;29:573-85.
- Davalos DB, Yadon CA, Tregellas HC. Untreated prenatal maternal depression and the potential risks to offspring: a review. *Arch Womens Ment Health* 2012;15:1-14.
- Davanzo R, Copertino M, De Cunto A, et al. Antidepressant drugs and breastfeeding: a review of the literature. *Breastfeed Med* 2011;6:89-98.
- Davis LL, Frazier E, Husain MM, et al. Substance use disorder comorbidity in major depressive disorder: a confirmatory analysis of the STAR*D cohort. *Am J Addict* 2006;15:278-85.
- de Maat S, Dekker J, Schoevers R, et al. Short psychodynamic supportive psychotherapy, antidepressants, and their combination in the treatment of major depression: a mega-analysis based on three randomized clinical trials. *Depress Anxiety* 2008;25:565-74.
- Delgado PL, Price LH, Charney DS, Heninger GR. Efficacy of fluvoxamine in treatment-refractory depression. *J Affect Disord* 1998;15:55-60.
- Dickinson LM, Rost K, Nutting PA, et al. RCT of a care manager intervention for major depression in primary care: 2-year costs for patients with physical vs psychological complaints. *Ann Fam Med* 2005;3:15-22.
- Dieserud G, Røysamb E, Ekeberg O, Kraft P. Toward an integrative model of suicide attempt: a cognitive psychological approach. *Suicide Life Threat Behav* 2001;31:153-68.
- Dietz PM, Williams SB, Callaghan WM, et al. Clinically identified maternal depression before, during, and after pregnancies ending in live births. *Am J Psychiatry* 2007;164:1515-20.
- Dimidjian S, Hollon SD, Dobson KS, et al. Randomized trial of behavioral activation, cognitive therapy, and antidepressant medication in the acute treatment of adults with major depression. *J Consult Clin Psychol* 2006;74:658-70.
- Dimitriou EC, Dimitriou CE. Buspirone augmentation of antidepressant therapy. *J Clin Psychopharmacol* 1998;18:465-69.
- Dobson KS, Hollon SD, Dimidjian S, et al. Randomized trial of behavioral activation, cognitive therapy, and antidepressant medication in the prevention of relapse and recurrence in major depression. *J Consult Clin Psychol* 2008;76:468-77.
- Donnelly PL. The use of the patient health questionnaire-9 Korean version (PHQ-9K) to screen for depressive disorders among Korean Americans. *J Transcult Nurs* 2007;18:324-30.
- Dørheim SK, Bondevik GT, Eberhard-Gran M, Bjorvatn B. Sleep and depression in postpartum women: a population-based study. *Sleep* 2009;32:847-55.
- Driessen E, Cuijpers P, Hollon SD, Dekker JJ. Does pretreatment severity moderate the efficacy of psychological treatment of adult outpatient depression? A meta-analysis. *J Consult Clin Psychol* 2010;78:668-80.
- Duffy FF, Chung H, Trivedi M, et al. Systematic use of patient-rated depression severity monitoring: is it helpful and feasible in clinical psychiatry? *Psychiatric Serv* 2008;59:1148-54.
- Dunlop BW, Crits-Christoph P, Evans DL, et al. Coadministration of modafinil and a selective serotonin reuptake inhibitor from the initiation of treatment of major depressive disorder with fatigue and sleepiness: a double-blind, placebo-controlled study. *J Clin Psychopharmacol* 2007;27:614-19.
- Dunn AL, Madhukar HT, Kampert JB, et al. Exercise treatment for depression: efficacy and dose response. *Am J Prev Med* 2005;28:1-8.

References

- Egede LE. Effect of comorbid chronic diseases on prevalence and odds of depression in adults with diabetes. *Psychosomatic Med* 2005;67:46-51.
- Ekers D, Murphy R, Archer J, et al. Nurse-delivered collaborative care for depression and long-term physical conditions: a systematic review and meta-analysis. *J Affect Disord* 2013;149:14-22.
- Ekers D, Richards D, Gilbody S. A meta-analysis of randomized trials of behavioural treatment of depression. *Psychol Med* 2008;38:611-23.
- Eng JJ, Reime B. Exercise for depressive symptoms in stroke patients: a systematic review and meta-analysis. *Clin Rehabil* 2014;28:731-39.
- Epperson CN, Terman M, Terman JS, et al. Randomized clinical trial of bright light therapy for antepartum depression: preliminary findings. *J Clin Psychiatry* 2004;65:421-25.
- Evans RW, Tepper SJ, Shapiro RE, et al. The FDA alert on serotonin syndrome with use of triptans combined with selective serotonin reuptake inhibitors or selective serotonin-norepinephrine reuptake inhibitors: American headache society position paper. *Headache* 2010;50:1089-99.
- Fava M, Rosenbaum JF, McGrath PJ, et al. Lithium and tricyclic augmentation of fluoxetine treatment for resistant major depression: a double-blind, controlled study. *Am J Psychiatry* 1994;151:1372-74.
- Fava M, Thase ME, DeBattista C. A multicenter, placebo-controlled study of modafinil augmentation in partial responders to selective serotonin reuptake inhibitors with persistent fatigue and sleepiness. *J Clin Psychiatry* 2005;66:85-93.
- Ferreira E, Carceller AM, Agogué C, et al. Effects of selective serotonin reuptake inhibitors and venlafaxine during pregnancy in term and preterm neonates. *Pediatrics* 2007;119:52-59.
- Fiellin DA, Carrington Reid M, O'Connor PG. Screening for alcohol problems in primary care: a systematic review. *Arch Intern Med* 2000;160:1977-89.
- Flint AJ, Rifat SL. Maintenance treatment for recurrent depression in late life. *Am J Geriatr Psychiatry* 2000;8:112-16.
- Flückiger C, Del Re AC, Munder T, et al. Enduring effects of evidence-based psychotherapies in acute depression and anxiety disorders versus treatment as usual at follow-up: a longitudinal meta-analysis. *Clin Psychol Rev* 2014;34:367-75.
- Flynn HA, Sexton M, Ratliff S, et al. Comparative performance of the Edinburgh postnatal depression scale and the patient health questionnaire-9 in pregnant and postpartum women seeking psychiatric services. *Psychiatry Res* 2011;187:130-34.
- Fond G, Loundou A, Rabu C, et al. Ketamine administration in depressive disorders: a systematic review and meta-analysis. *Psychopharmacology* 2014;231:3663-76.
- Fortney JC, Pyne JM, Mouden SB, et al. Practice-based versus telemedicine-based collaborative care for depression in rural federally qualified health centers: a pragmatic randomized comparative effectiveness trial. *Am J Psychiatry* 2013;170:414-25.
- Fournier JC, DeRubeis RJ, Hollon SD, et al. Antidepressant drug effects and depression severity: a patient-level meta-analysis. *JAMA* 2010;303:47-53.
- Fraguas R Jr, Iosifescu DV, Alpert J, et al. Major depressive disorder and comorbid cardiac disease: is there a depressive subtype with greater cardiovascular morbidity? results from the STAR*D study. *Psychosomatics* 2007;48:418-25.
- Frank E, Kupfer DJ, Perel JM, et al. Comparison of full-dose versus half-dose pharmacotherapy in the maintenance treatment of recurrent depression. *J Affect Disord* 1993;27:139-45.

References

- Frasure-Smith N, Lespérance F, Talajic M. Depression in 18-month prognosis after myocardial infarction. *Circulation* 1995;91:999-1005.
- Freeman MP, Davis M, Sinha P, et al. Omega-3 fatty acids and supportive psychotherapy for perinatal depression: a randomized placebo-controlled study. *J Affect Disord* 2008;110:142-48.
- Freeman MP, Fava M, Lake J, et al. Complementary and alternative medicine in major depressive disorder: the American psychiatric association task force report. *J Clin Psychiatry* 2010;71:669-81.
- Frye MA. Bipolar disorder – a focus on depression. *N Engl J Med* 2011;364:51-59.
- Ganzini L, Smith DM, Fenn DS, Lee MA. Depression and mortality in medically ill older patients. *J Am Geriatr Soc* 1997;45:307-12.
- Garakani A, Win T, Virk S, et al. Comorbidity of irritable bowel syndrome in psychiatric patients: a review. *Am J of Therapeutics* 2003;10:61-67.
- Garfield S, Francis SA, Smith FJ. Building concordant relationships with patients starting antidepressant medication. *Patient Educ Couns* 2004;55:241-46.
- Garrard JM. Patient outcomes associated with antidepressant drugs. Agency for Healthcare Research and Quality (AHRQ). AHRQ 2001-64. April 2001.
- Gartlehner G, Gaynes BN, Hansen RA, et al. Comparative benefits and harms of second-generation antidepressants: background paper for the American college of physicians. *Ann Intern Med* 2008;148:734-50.
- Gastó C, Navarro V, Marcos T, et al. Single-blind comparison of venlafaxine and nortriptyline in elderly major depression. *J Clin Psychopharmacol* 2003;23:21-26.
- Gaynes BN, DeVaugh-Geiss J, Weir S, et al. Feasibility and diagnostic validity of the M-3 checklist: a brief, self-rated screen for depressive, bipolar, anxiety, and post-traumatic stress disorders in primary care. *Ann Fam Med* 2010;8:160-69.
- Gaynes BN, Gavin N, Meltzer-Brody S, et al. Perinatal depression: prevalence, screening accuracy, and screening outcomes. Summary, evidence report/technology assessment no. 119. (Prepared by the RTI-University of North Carolina evidence-based practice center under contract no. 290-02-0016.) AHRQ publication no. 05-E006-1. Rockville, MD: Agency for Healthcare Research and Quality. February 2005.
- Geddes JR, Carney SM, Davies C, et al. Relapse prevention with antidepressant drug treatment in depressive disorders: a systematic review. *Lancet* 2003;361:653-61.
- George MS, Lisanby SH, Avery D, et al. Daily left prefrontal transcranial magnetic stimulation therapy for major depressive disorder: a sham-controlled randomized trial. *Arch Gen Psychiatry* 2010;67:507-16.
- George MS, Rush AJ, Marangell LB, et al. A one-year comparison of vagus nerve stimulation with treatment as usual for treatment-resistant depression. *Biol Psychiatry* 2005;58:364-73.
- Gilbody S, Bower P, Fletcher J, et al. Collaborative care for depression: a cumulative meta-analysis and review of longer-term outcomes. *Arch Intern Med* 2006;166:2314-21.
- Gilbody S, Richards D, Brealey S, Hewitt C. Screening for depression in medical settings with the patient health questionnaire (PHQ): a diagnostic meta-analysis. *J Gen Intern Med* 2007;22:1596-602.
- Gilchrist G, Hegarty K, Chondros P, et al. The association between intimate partner violence, alcohol and depression in family practice. *BMC Fam Pract* 2010;11:72.
- Gillman PK. A review of serotonin toxicity data: implications for the mechanisms of antidepressant drug action. *Biol Psychiatry* 2006;59:1046-51.
- Gillman PK. Triptans, serotonin agonists, and serotonin syndrome (serotonin toxicity): a review. *Headache* 2010;50:264-72.

References

- Gjerdingen D, Crow S, McGovern P, et al. Postpartum depression screening at well-child visits: validity of a 2-question screen and the PHQ-9. *Ann Fam Med* 2009;7:63-70.
- Gjerdingen DK, Yawn BP. Postpartum depression screening: importance, methods, barriers, and recommendations for practice. *J Am Board Fam Med* 2007;20:280-88.
- Gold KJ, Singh V, Marcus SM, Palladino CL. Mental health, substance use and intimate partner problems among pregnant and postpartum suicide victims in the national violent death reporting system. *Gen Hosp Psychiatry* 2012;34:139-45.
- Golden RN, Gaynes BN, Ekstrom RD, et al. The efficacy of light therapy in the treatment of mood disorders: a review and meta-analysis of the evidence. *Am J Psychiatry* 2005;162:656-62.
- Goodwin FK, Jamison KR. Clinical depression and diagnosis. In Manic-Depressive Illness: Bipolar Disorders and Recurrent Depression. 2nd Edition. Chapter 1. New York: Oxford University Press. 2007.
- Goyal D, Gay C, Lee K. Fragmented maternal sleep is more strongly correlated with depressive symptoms than infant temperament at three months postpartum. *Arch Womens Ment Health* 2009;12:229-37.
- Greden JF. Antidepressant maintenance medications: when to discontinue and how to stop. *J Clin Psychiatry* 1993;54:39-47.
- Green RC, Cupples A, Kurz A, et al. Depression as a risk factor for alzheimer disease: the MIRAGE study. *Arch Neurol* 2003;60:753-59.
- Gregory RJ, Canning SS, Lee TW, Wise JC. Cognitive bibliotherapy for depression: a meta-analysis. *Prof Psychology Res and Prac* 2004;35:275-80.
- Guidi J, Fava GA, Fava M, Papakostas GI. Efficacy of the sequential integration of psychotherapy and pharmacotherapy in major depressive disorder: a preliminary meta-analysis. *Psychol Med* 2011;41:321-31.
- Haby MM, Donnelly M, Corry J, Vos T. Cognitive behavioural therapy for depression, panic disorder and generalized anxiety disorder: a meta-regression of factors that may predict outcome. *Aust NZ J Psychiatry* 2006;40:9-19.
- Hamann J, Cohen R, Luecht S, et al. Do patients with schizophrenia wish to be involved in decisions about their medical treatment? *Am J Psychiatry* 2005;162:2382-84.
- Han C, Jo SA, Kwak J, et al. Validation of the patient health questionnaire-9 Korean version in the elderly population: the Ansan geriatric study. *Compr Psychiatry* 2008;49:218-23.
- Hausenblas HA, Saha D, Dubyak PJ, Anton SD. Saffron (*Crocus sativus L.*) and major depressive disorder: a meta-analysis of randomized clinical trials. *J Integr Med* 2013;11:377-83.
- Hawkins EJ, Kivlahan DR, Williams EC, et al. Examining quality issues in alcohol misuse screening. *Subst Abus* 2007;28:53-65.
- Hedayati SS, Minhajuddin AT, Afshar M, et al. Association between major depressive episodes in patients with chronic kidney disease and initiation of dialysis, hospitalization, or death. *JAMA* 2010;303:1946-53.
- Hirschfeld RMA. Clinical importance of long-term antidepressant treatment. *Br J Psychiatry Suppl* 2001;179:S4-S8.
- Hirschfeld RMA, Williams JBW, Spitzer RL, et al. Development and validation of a screening instrument for bipolar spectrum disorder: the mood disorder questionnaire. *Am J Psychiatry* 2000;157:1873-75.
- Hofmann SG, Sawyer AT, Witt AA, Oh D. The effect of mindfulness-based therapy on anxiety and

References

- depression: a meta-analytic review. *J Consult Clin Psychol* 2010;78:169-83.
- Hollon SD, DeRubeis RJ, Fawcett J, et al. Effect of cognitive therapy with antidepressant medications vs antidepressants alone on the rate of recovery in major depressive disorder: a randomized clinical trial. *JAMA Psychiatry* 2014;71:1157-64.
- Huang FY, Chung H, Kroenke K, et al. Using the patient health questionnaire-9 to measure depression among racially and ethnically diverse primary care patients. *J Gen Intern Med* 2006;21:547-52.
- Huijbregts KML, de Jong FJ, van Marwijk HWJ, et al. A target-driven collaborative care model for major depressive disorder is effective in primary care in the Netherlands: a randomized clinical trial from the depression initiative. *J Affect Dis* 2013;146:328-37.
- Hunkeler EM, Katon W, Tang L, et al. Long term outcomes from the IMPACT randomised trial for depressed elderly patients in primary care. *BMJ* 2006;332:259-63.
- Hunkeler EM, Meresman JF, Hargreaves WA, et al. Efficacy of nurse telehealth care and peer support in augmenting treatment of depression in primary care. *Arch Fam Med* 2000;9:700-08.
- Ialongo N, McCreary BK, Pearson JL, et al. Major depressive disorder in a population of urban, African-American young adults: prevalence, correlates, comorbidity and unmet mental health service need. *J Affect Disord* 2004;79:127-36.
- Imel ZE, Malterer MB, McKay KM, Wampold BE. A meta-analysis of psychotherapy and medication in unipolar depression and dysthymia. *J Affect Disord* 2008;110:197-206.
- Jakobsen JC, Hansen JL, Simonsen S, et al. Effects of cognitive therapy versus interpersonal psychotherapy in patients with major depressive disorder: a systematic review of randomized clinical trials with meta-analyses and trial sequential analyses. *Psychol Med* 2012a;42:1343-57.
- Jakobsen JC, Hansen JL, Simonsen E, Gluud C. The effect of adding psychodynamic therapy to antidepressants in patients with major depressive disorder. A systematic review of randomized clinical trials with meta-analyses and trial sequential analyses. *J Affect Disord* 2012b;137:4-14.
- Jakobsen JC, Hansen JL, Simonsen E, Gluud C. The effect of interpersonal psychotherapy and other psychodynamic therapies versus 'treatment as usual' in patients with major depressive disorder. *PLoS One* 2011a;6:e19044.
- Jakobsen JC, Hansen JL, Storebø OJ, et al. The effects of cognitive therapy versus 'no intervention' for major depressive disorder. *PLoS One* 2011b;6:e28299.
- Jakobsen JC, Lindschou Hansen J, Storebø OJ, et al. The effects of cognitive therapy versus 'treatment as usual' in patients with major depressive disorder. *PLoS One* 2011c;6:e22890.
- Jakubovski E, Varigonda AL, Freemantle N, et al. Systematic review and meta-analysis: dose-response relationship of selective serotonin reuptake inhibitors in major depressive disorder. *Am J Psychiatry* 2016;173:174-83.
- Janicak PG, Nahas Z, Lisanby SH, et al. Durability of clinical benefit with transcranial magnetic stimulation (TMS) in the treatment of pharmacoresistant major depression: assessment of relapse during a 6-month, multisite, open-label study. *Brain Stimul* 2010;3:187-99.
- Jeeva F, Dickens C, Coventry P. Is treatment of depression cost-effective in people with diabetes? A systematic review of the economic evidence. *Int J Technol Assess Health Care* 2013;29:384-91.
- Jiang W, Davidson JRT. Antidepressant therapy in patients with ischemic heart disease. *Am Heart J* 2005;150:871-81.
- Joffe RT, Singer W, Levitt AJ, et al. A placebo-controlled comparison of lithium and triiodothyronine augmentation of tricyclic antidepressants in unipolar refractory depression. *Arch Gen Psychiatry*

References

1993;50:387-93.

Johnston BJ. The role of patient experience and its influence on adherence to antidepressant treatment. *J Psychosoc Nurs Ment Health Serv* 2013;51:29-37.

Jorm AF, Christensen H, Griffiths KM, Rodgers B. Effectiveness of complementary and self-help treatments for depression. *Med J Aust* 2002;176:S84-S96.

Judd LL, Akiskal HS, Schettler PJ, et al. The long-term natural history of the weekly symptomatic status of bipolar I disorder. *Arch Gen Psychiatry* 2002;59:530-37.

Karasz A. Cultural differences in conceptual models of depression. *Social Science & Medicine* 2005;60:1625-35.

Katon W, Robinson P, Von Korff M, et al. A multifaceted intervention to improve treatment of depression in primary care. *Arch Gen Psychiatry* 1996;53:924-32.

Katon W, Von Korff M, Lin E, et al. Stepped collaborative care for primary care patients with persistent symptoms of depression: a randomized trial. *Arch Gen Psychiatry* 1999;56:1109-15.

Katon W, Von Korff M, Ciechanowski P, et al. Behavioral and clinical factors associated with depression among individuals with diabetes. *Diabetes Care* 2004a;27:914-20.

Katon WJ, Lin EHB, Russo J, et al. Cardiac risk factors in patients with diabetes mellitus and major depression. *J Gen Intern Med* 2004b;19:1192-99.

Katon WJ, Seelig M. Population-based care of depression: team care approaches to improving outcomes. *J Occup Environ Med* 2008;50:459-67.

Katon WJ, Schoenbaum M, Fan M, et al. Cost-effectiveness of improving primary care treatment of late-life depression. *Arch Gen Psychiatry* 2005;62:1313-20.

Katona C, Abou-Saleh MT, Harrison DA, et al. Placebo-controlled trial of lithium augmentation of fluoxetine and lofepramine. *Br J Psychiatry* 1995;166:80-86.

Keller MB. Issues in treatment-resistant depression. *J Clin Psychiatry* 2005;66:5-12.

Keller MB, Kocsis JH, Thase ME, et al. Maintenance phase efficacy of sertraline for chronic depression: a randomized controlled trial. *JAMA* 1998;280:1665-72.

Keller MB, McCullough JP, Klein DN, et al. A comparison of nefazodone, the cognitive behavioral-analysis system of psychotherapy, and their combination for the treatment of chronic depression. *N Engl J Med* 2000;342:1462-70.

Kellner CH, Knapp RG, Petrides G, et al. Continuation electroconvulsive therapy vs pharmacotherapy for relapse prevention in major depression: a multisite study from the consortium for research in electroconvulsive therapy (CORE). *Arch Gen Psychiatry* 2006;63:1337-44.

Kennedy SH, Giacobbe P, Rizvi SJ, et al. Deep brain stimulation for treatment-resistant depression: follow-up after 3 to 6 years. *Am J Psychiatry* 2011;168:502-10.

Kessler RC, Chiu WT, Demler O, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the national comorbidity survey replication. *Arch Gen Psychiatry* 2005;62:617-27.

Kieler H, Artama M, Engeland A, et al. Selective serotonin reuptake inhibitors during pregnancy and risk of persistent pulmonary hypertension in the newborn: population based cohort study from the five Nordic countries. *BMJ* 2011;344:d8012.

Kim KH, Lee SM, Paik JW, Kim NS. The effects of continuous antidepressant treatment during the first 6 months on relapse or recurrence of depression. *J Affect Disord* 2011;132:121-29.

References

- Kirmayer LJ. Cultural variations in the clinical presentation of depression and anxiety: implications for diagnosis and treatment. *J Clin Psychiatry* 2001;62:22-28.
- Kleinman A. Culture and depression. *N Engl J Med* 2004;351:951-53.
- Klinkman MS. The role of algorithms in the detection and treatment of depression in primary care. *J Clin Psychiatry* 2003;64:19-23.
- Kocsis JH, Gelenberg AJ, Rothbaum BO, et al. Cognitive behavioral analysis system of psychotherapy and brief supportive psychotherapy for augmentation of antidepressant nonresponse in chronic depression: the REVAMP trial. *Arch Gen Psychiatry* 2009a;66:1178-88.
- Kocsis JH, Leon AC, Markowitz JC, et al. Patient preference as a moderator of outcome for chronic forms of major depressive disorder treated with nefazodone, cognitive behavioral analysis system of psychotherapy, or their combination. *J Clin Psychiatry* 2009b;70:354-61.
- Kok RM, Heeren TJ, Nolen WA. Continuing treatment of depression in the elderly: a systematic review and meta-analysis of double-blinded randomized controlled trials with antidepressants. *Am J Geriatr Psychiatry* 2011;19:249-55.
- Kok RM, Nolen WA, Heeren TJ. Efficacy of treatment in older depressed patients: a systematic review and meta-analysis of double-blind randomized controlled trials with antidepressants. *J Affect Disord* 2012;141:103-15.
- Kozhimannil KB, Pereira MA, Harlow BL. Association between diabetes and perinatal depression among low-income mothers. *JAMA* 2009;301:842-47.
- Kraft JB, Slager SL, McGrath PJ, Hamilton SP. Sequence analysis of the serotonin transporter and associations with antidepressant response. *Biol Psychiatry* 2005;58:374-81.
- Kriston L, von Wolff A, Westphal A, et al. Efficacy and acceptability of acute treatments for persistent depressive disorder: a network meta-analysis. *Depress Anxiety* 2014;31:621-30.
- Kroenke K, Spitzer RL, Williams JBW, Löwe B. The patient health questionnaire somatic, anxiety, and depressive symptom scales: a systematic review. *Gen Hosp Psychiatry* 2010;32:345-59.
- Kroenke K, Spitzer RL, Williams JBW. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;16:606-13.
- Kroenke K, Wood DR, Mangelsdorff AD, et al. Chronic fatigue in primary care: prevalence, patient characteristics, and outcome. *JAMA* 1988;260:929-34.
- Krupnick JL, Sotsky SM, Simmens S, et al. The role of the therapeutic alliance in psychotherapy and pharmacotherapy outcome: findings in the national institute of mental health treatment of depression collaborative research program. *J Consult Clin Psychol* 1996;64:532-39.
- Kuyken W, Hayes R, Barrett B, et al. Effectiveness and cost-effectiveness of mindfulness-based cognitive therapy compared with maintenance antidepressant treatment in the prevention of depressive relapse or recurrence (PREVENT): a randomised controlled trial. *Lancet* 2015;386:63-73.
- Lancaster CA, Gold KJ, Flynn HA, et al. Risk factors for depressive symptoms during pregnancy: a systematic review. *Am J Obstet Gynecol* 2010;202:5-14.
- Lavretsky H, Reinlieb M, St. Cyr N, et al. Citalopram, methylphenidate, or their combination in geriatric depression: a randomized, double-blind, placebo-controlled trial. *Am J Psychiatry* 2015;172:561-69.
- Leichsenring F, Rabung S, Leibing E. The efficacy of short-term psychodynamic psychotherapy in

References

- specific psychiatric disorders: a meta-analysis. *Arch Gen Psychiatry* 2004;61:1208-16.
- Lépine JP, Briley M. The epidemiology of pain in depression. *Hum Psychopharmacol* 2004;19:S3-S7.
- Leppämäki SJ, Partonen TT, Hurme J, et al. Randomized trial of the efficacy of bright-light exposure and aerobic exercise on depressive symptoms and serum lipids. *J Clin Psychiatry* 2002;63:316-21.
- Levine S, Unützer J, Yip JY, et al. Physicians' satisfaction with a collaborative disease management program for late-life depression in primary care. *Gen Hosp Psychiatry* 2005;27:383-91.
- [Levkovitz Y, Tedeschini E, Papakostas GI](#). Efficacy of antidepressants for dysthymia: a meta-analysis of placebo-controlled randomized trials. *J Clin Psychiatry* 2011;72:509-14.
- Li D, Liu L, Odouli R. Presence of depressive symptoms during early pregnancy and the risk of preterm delivery: a prospective cohort study. *Hum Reprod* 2009;24:146-53.
- [Li G, Mbuagbaw L, Samaan Z, et al](#). Efficacy of vitamin D supplementation in depression in adults: a systematic review. *J Clin Endocrinol Metab* 2014;99:757-67.
- Lichtman JH, Bigger Jr JT, Blumenthal JA, et al. Depression and coronary heart disease: recommendations for screening, referral, and treatment: a science advisory from the American heart association prevention committee of the council on cardiovascular nursing, council on clinical cardiology, council on epidemiology and prevention, and interdisciplinary council on quality of care and outcomes research: endorsed by the American psychiatric association. *Circulation* 2008;118:1768-75.
- Linde K, Berner MM, Kriston L. St John's wort for major depression. *Cochrane Database Syst Rev* 2008:CD000448.
- Lo Sasso AT, Rost K, Beck A. Modeling the impact of enhanced depression treatment on workplace functioning and costs: a cost-benefit approach. *Med Care* 2006;44:352-58. (Cost Effectiveness Analysis)
- Loh A, Leonhart R, Wills CE, et al. The impact of patient participation on adherence and clinical outcome in primary care of depression. *Patient Educ Couns* 2007;65:69-78.
- Lotrakul M, Sumrithe S, Saipanish R. Reliability and validity of the Thai version of the PHQ-9. *BMC Psychiatry* 2008; 8:46.
- [Louie L](#). The effectiveness of yoga for depression: a critical literature review. *Issues Ment Health Nurs* 2014;35:265-76.
- Löwe B, Unützer J, Callahan CM, et al. Monitoring depression treatment outcomes with the patient health questionnaire-9. *Med Care* 2004;42:1194-1201.
- Lozano AM, Mayberg HS, Giacobbe P, et al. Subcallosal cingulate gyrus deep brain stimulation for treatment-resistant depression. *Biol Psychiatry* 2008;64:461-67.
- Ludman EJ, Katon W, Russo J, et al. Depression and diabetes symptom burden. *Gen Hosp Psychiatry* 2004;26:430-36.
- [MacPherson H, Richmond S, Bland M, et al](#). Acupuncture and counselling for depression in primary care: a randomised controlled trial. *PLoS Med* 2013;10:e1001518.
- Maes M, Libbrecht I, van Hunsel F, et al. Pindolol and mianserin augment the antidepressant activity of fluoxetine in hospitalized major depressed patients, including those with treatment resistance. *J Clin Psychopharmacol* 1999;19:177-82.
- [Major Depressive Disorder Working Group of the Psychiatric GWAS Consortium, Ripke S, Wray NR, et al](#). A mega-analysis of genome-wide association studies for major depressive disorder. *Mol Psychiatry* 2013;18:497-511.
- Malone Jr DA, Dougherty DD, Rezai AR, et al. Deep brain stimulation of the ventral capsule/ventral

References

- striatum for treatment-resistant depression. *Biol Psychiatry* 2009;65:267-75.
- Maneeton N, Maneeton B, Srisurapanont M, Martin SD. Quetiapine monotherapy in acute phase for major depressive disorder: a meta-analysis of randomized, placebo-controlled trials. *BMC Psychiatry* 2012;12:160.
- Manicavasagar V, Parker G, Perich T. Mindfulness-based cognitive therapy vs cognitive behaviour therapy as a treatment for non-melancholic depression. *J Affect Disord* 2011;130:138-44.
- Manu P, Matthews DA, Lane TJ. The mental health of patients with a chief complaint of chronic fatigue: a prospective evaluation and follow-up. *Arch Intern Med* 1988;148:2213-17.
- Marcus RN, McQuade RD, Carson WH, et al. The efficacy and safety of aripiprazole as adjunctive therapy in major depressive disorder: a second multicenter, randomized, double-blind, placebo-controlled study. *J Clin Psychopharmacol* 2008;28:156-65.
- Markowitz JC, Kocsis JH, Bleiberg KL, et al. A comparative trial of psychotherapy and pharmacotherapy for "pure" dysthymic patients. *J Affect Disord* 2005;89:167-75.
- Marshall RD, Liebowitz MR. Paroxetine/bupropion combination treatment for refractory depression. *J Clin Psychopharmacol* 1996;16:80-81.
- Mayberg HS, Lozano AM, Voon V, et al. Deep brain stimulation for treatment-resistant depression. *Neuron* 2005;45:651-60.
- Mazzucchelli T, Kane R, Rees C. Behavioral activation treatments for depression in adults: a meta-analysis and review. *Cin Psychol Sci Prac* 2009;16:383-411.
- McCarney RW, Schulz J, Grey AR. Effectiveness of mindfulness-based therapies in reducing symptoms of depression: a meta-analysis. *Eur J Psychotherapy Couns* 2012;14:279-99.
- McGrath PJ, Stewart JW, Nunes EN, et al. Treatment response of depressed outpatients unresponsive to both a tricyclic and a monoamine oxidase inhibitor antidepressant. *J Clin Psychiatry* 1994;55:336-39.
- McGrath PJ, Stewart JW, Nunes EV, et al. A double-blind crossover trial of imipramine and phenelzine for outpatients with treatment-refractory depression. *Am J Psychiatry* 1993;150:118-23.
- Meehan J, Kapur N, Hunt IM, et al. Suicide in mental health in-patients and within 3 months of discharge: national clinical survey. *Br J Psychiatry* 2006;188:129-34.
- Melfi CA, Chawla AJ, Croghan TW, et al. The effects of adherence to antidepressant treatment guidelines on relapse and recurrence of depression. *Arch Gen Psychiatry* 1998;55:1128-32.
- Menchetti M, Rucci P, Bortolotti B, et al. Moderators of remission with interpersonal counselling or drug treatment in primary care patients with depression: randomised controlled trial. *Br J Psychiatry* 2014;204:144-50.
- Merikangas KR, Ames M, Cui L, et al. The impact of comorbidity of mental and physical conditions on role disability in the US adult household population. *Arch Gen Psychiatry* 2007;64:1180-88.
- Mian AI. Depression in pregnancy and the postpartum period: balancing adverse effects of untreated illness and treatment risks. *J Psychiatr Pract* 2005;11:389-96.
- Midanik LT, Zahnd EG, Klein D. Alcohol and drug CAGE screeners for pregnant, low-income women: the California perinatal needs assessment. *Alcohol Clin Exp Res* 1998;22:121-25.
- Milgrom J, Gemmill AW. Screening for perinatal depression. *Best Pract Res Clin Obstet Gynaecol* 2014;28:13-23.
- Miranda J, Schoenbaum M, Sherbourne C, et al. Effects of primary care depression treatment on minority patients' clinical status and employment. *Arch Gen Psychiatry* 2004;61:827-34.

References

- Mitchell AJ, Subramaniam H. Prognosis of depression in old age compared to middle age: a systematic review of comparative studies. *Am J Psychiatry* 2005;162:1588-1601.
- Mojtabai R, Olfson M. National patterns in antidepressant treatment by psychiatrists and general medical providers: results from the national comorbidity survey replication. *J Clin Psychiatry* 2008;69:1064-74.
- Morgan AJ, Jorm AF. Self-help interventions for depressive disorders and depressive symptoms: a systematic review. *Ann Gen Psychiatry* 2008;7:13.
- Moussavi S, Chatterji S, Verdes E, et al. Depression, chronic diseases, and decrements in health: results from the World Health Surveys. *Lancet* 2007;370:851-58.
- Muñoz RF, Mendelson T. Toward evidence-based interventions for diverse populations: the San Francisco general hospital prevention and treatment manuals. *J Consulting Clin Psychol* 2005;73:790-99.
- Murrough JW, Iosifescu DV, Chang LC, et al. Antidepressant efficacy of ketamine in treatment-resistant major depression: a two-site randomized controlled trial. *Am J Psychiatry* 2013;170:1134-42.
- Nahas Z, Marangell LB, Husian MM, et al. Two-year outcome of vagus nerve stimulation (VNS) for treatment of major depression episodes. *J Clin Psychiatry* 2005;66:1097-1104.
- Narasimhan S, Lohoff FW. Pharmacogenetics of antidepressant drugs: current clinical practice and future directions. *Pharmacogenomics* 2012;13:441-64.
- National Institute for Clinical Excellence. Guidance on the use of electroconvulsive therapy. Technology Appraisal Guidance 59. London:England. April 2003.
- National Institute for Health and Care Excellence. Depression in adults. Available at: nice.org.uk/guidance/qs8. 2011.
- Nelson JC. Augmentation strategies in depression 2000. *J Clin Psychiatry* 2000;61:13-19.
- Nelson JC, Mazure CM, Bowers MB Jr, Jatlow PI. A preliminary, open study of the combination of fluoxetine and desipramine for rapid treatment of major depression. *Arch Gen Psychiatry* 1991;48:303-07.
- Nelson JC, Papakostas GI. Atypical antipsychotic augmentation in major depressive disorder: a meta-analysis of placebo-controlled randomized trials. *Am J Psychiatry* 2009;166:980-91.
- Nichita EC, Buckley PF. Informed consent and competency: doctor's dilemma on the consultation liaison service. *Psychiatry (Edgmont)* 2007;4:53-55.
- Nicholson A, Kuper H, Hemingway H. Depression as an aetiologic and prognostic factor in coronary heart disease: a meta-analysis of 6,362 events among 146,538 participants in 54 observational studies. *Eur Heart J* 2006;27:2763-74.
- Nierenberg AA, Fava M, Trivedi MH, et al. A comparison of lithium and T₃ augmentation following two failed medication treatments for depression: a STAR*D report. *Am J Psychiatry* 2006;163:1519-30.
- Nierenberg AA, McColl RD. Management options for refractory depression. *Am J Med* 1996;101:45S-52S.
- Nimalasuriya K, Compton MT, Guillory VJ. Screening adults for depression in primary care: a position statement of the American college of preventive medicine. *J Fam Pract* 2009;58:535-38.
- NIMH/NIH Consensus Development Conference Statement. Mood disorders: pharmacologic prevention of recurrences. *Am J Psychiatry* 1985;142:469-76.
- Ninan PT, Hassman HA, Glass SJ, McManus FC. Adjunctive modafinil at initiation of treatment with a selective serotonin reuptake inhibitor enhances the degree and onset of therapeutic effects in patients with major depressive disorder and fatigue. *J Clin Psychiatry* 2004;65:414-20.

References

- Oberlander TF, Warburton W, Misri S, et al. Neonatal outcomes after prenatal exposure to selective serotonin reuptake inhibitor antidepressants and maternal depression using population-based linked health data. *Arch Gen Psychiatry* 2006;63:898-906.
- O'Connor AM, Bennett C, Stacey D, et al. Do patient decision aids meet effectiveness criteria of the international patient decision aid standards collaboration? A systematic review and meta-analysis. *Med Decis Making* 2007;27:554-74.
- O'Connor EA, Rossom RC, Henninger M, et al. Screening for depression in adults: an updated systematic evidence review for the U.S. preventive services task force. AHRQ Publication No. 14-05208-EF-1. January 2016.
- Oestergaard S, Møldrup C. Optimal duration of combined psychotherapy and pharmacotherapy for patients with moderate and severe depression: a meta-analysis. *J Affect Disord* 2011;131:24-36.
- O'Hara MW. Postpartum depression: what we know. *J Clin Psychol* 2009;65:1258-69.
- O'Hara MW, Stuart S, Gorman LL, Wenzel A. Efficacy of interpersonal psychotherapy for postpartum depression. *Arch Gen Psychiatry* 2000;57:1039-45.
- O'Hara MW, Wisner KL. Perinatal mental illness: definition, description and aetiology. *Best Pract Res Clin Obstet Gynaecol* 2014;28:3-12.
- Okumura Y, Ichikura K. Efficacy and acceptability of group cognitive behavioral therapy for depression: a systematic review and meta-analysis. *J Affect Disord* 2014;164:155-64.
- Onyike CU, Crum RM, Lee HB, et al. Is obesity associated with major depression?: results from the third national health and nutrition examination survey. *Am J Epidemiol* 2003;158:1139-47.
- Oquendo MA, Friend JM, Halberstam B, et al. Association of comorbid posttraumatic stress disorder and major depression with greater risk for suicidal behavior. *Am J Psychiatry* 2003;160:580-82.
- O'Reardon JP, Solvason HB, Janicak PG, et al. Efficacy and safety of transcranial magnetic stimulation in the acute treatment of major depression: a multisite randomized controlled trial. *Biol Psychiatry* 2007; 62:1208-16.
- Oren DA, Wisner KL, Spinelli M, et al. An open trial of morning light therapy for treatment of antepartum depression. *Am J Psychiatry* 2002;159:666-69.
- Ownby RL, Crocco E, Acevedo A, et al. Depression and risk for alzheimer disease: systematic review, meta-analysis, and metaregression analysis. *Arch Gen Psychiatry* 2006;63:530-38.
- Oxman TE, Dietrich AJ, Williams JW Jr, Kroenke K. A three-component model for reengineering systems for the treatment of depression in primary care. *Psychosomatics* 2002;43:441-50.
- Pan A, Sun Q, Okereke OI, et al. Depression and risk of stroke morbidity and mortality: a meta-analysis and systematic review. *JAMA* 2011;306:1241-49.
- Papakostas GI, Shelton RC, Smith J, Fava M. Augmentation of antidepressants with atypical antipsychotic medications for treatment-resistant major depressive disorder: a meta-analysis. *J Clin Psychiatry* 2007;68:826-31.
- Paulson JF, Bazemore SD. Prenatal and postpartum depression in fathers and its association with maternal depression: a meta-analysis. *JAMA* 2010;303:1961-69.
- Paykel ES, Ramana R, Cooper Z, et al. Residual symptoms after partial remission: an important outcome in depression. *Psychological Medicine* 1995;25:1171-80.
- Pearlstein T. Perinatal depression: treatment options and dilemmas. *J Psychiatry Neurosci* 2008;33:302-

References

18.

Peeters F, Huibers M, Roelofs J, et al. The clinical effectiveness of evidence-based interventions for depression: a pragmatic trial in routine practice. *J Affect Disord* 2013;145:349-55.

Perry PJ, Zeilmann C, Arndt S. Tricyclic antidepressant concentrations in plasma: an estimate of their sensitivity and specificity as a predictor of response. *J Clin Psychopharmacol* 1994;14:230-40.

Petrak F, Herpertz S, Albus C, et al. Study protocol of the diabetes and depression study (DAD): a multi-center randomized controlled trial to compare the efficacy of a diabetes-specific cognitive behavioral group therapy versus sertraline in patients with major depression and poorly controlled diabetes mellitus. *BMC Psychiatry* 2013;13:206.

Pettersson A, Boström KB, Gustavsson P, Ekselius L. Which instruments to support diagnosis of depression have sufficient accuracy? A systematic review. *Nord J Psychiatry* 2015;69:497-508.

Piet J, Hougaard E. The effect of mindfulness-based cognitive therapy for prevention of relapse in recurrent major depressive disorder: a systematic review and meta-analysis. *Clin Psychol Rev* 2011;31:1032-40.

Pinquart M, Duberstein PR, Lyness JM. Effects of psychotherapy and other behavioral interventions on clinically depressed older adults: a meta-analysis. *Aging Ment Health* 2007;11:645-57.

Pinto-Meza A, Serrano-Blanco A, Peñarrubia M, et al. Assessing depression in primary care with the PHQ-9: can it be carried out over the telephone? *J Gen Intern Med* 2005;20:738-42.

Pizzi C, Rutjes AW, Costa GM, et al. Meta-analysis of selective serotonin reuptake inhibitors in patients with depression and coronary heart disease. *Am J Cardiol* 2011;107:972-79.

Porcelli S, Fabbri C, Drago A, et al. Genetics and antidepressant: where we are. *Clin Neuropsychiatry* 2011;8:99-150.

Post RM. Transduction of psychosocial stress into the neurobiology of recurrent affective disorder. *Am J Psychiatry* 1992;149:999-1010.

Prasko J, Horacek J, Klaschka J, et al. Bright light therapy and/or imipramine for inpatients with recurrent non-seasonal depression. *Neuro Endocrinol Lett* 2002;23:109-13.

Preskorn SH. Ketamine: the hopes and the hurdles. *Biol Psychiatry* 2012;72:522-23.

Preskorn SH, Beber JH, Faul JC, Hirschfeld R. Serious adverse effects of combining fluoxetine and tricyclic antidepressants. *Am J Psychiatry* 1990;147:532.

Ravindran AV, Lam RW, Filteau MJ, et al. Canadian network for mood and anxiety treatments (CANMAT) Clinical guidelines for the management of major depressive disorder in adults. V. complementary and alternative medicine treatments. *J Affect Disord* 2009;117:S54-64.

Ray S, Nizamie SH, Akhtar S, et al. Efficacy of adjunctive high frequency repetitive transcranial magnetic stimulation of left prefrontal cortex in depression: a randomized sham controlled study. *J Affect Disord* 2011;128:153-39.

Ray S, Stowe ZN. The use of antidepressant medication in pregnancy. *Best Pract Res Clin Obstet Gynaecol* 2014;28:71-83.

Rethorst CD, Sunderajan P, Greer TL, et al. Does exercise improve self-reported sleep quality in non-remitted major depressive disorder? *Psychol Med* 2013;43:699-709.

Reynolds III CF. Treatment of depression in special populations. *J Clin Psychiatry* 1992;53:45-53.

Reynolds III CF, Dew MA, Pollock BG, et al. Maintenance treatment of major depression in old age. *N Engl J Med* 2006;354:1130-38.

References

Robinson M, Oakes TM, Raskin J, et al. Acute and long-term treatment of late-life major depressive disorder: duloxetine versus placebo. *Am J Geriatr Psychiatry* 2014;22:34-45.

Rost K, Nutting P, Smith JL, et al. Managing depression as a chronic disease: a randomised trial of ongoing treatment in primary care. *BMJ* 2002;325:934-37.

Rost K, Smith JL, Dickinson M, et al. The effect of improving primary care depression management on employee absenteeism and productivity: a randomized trial. *Med Care* 2004;42:1202-10.

Rugulies R. Depression as a predictor for coronary heart disease: a review and meta-analysis. *Am J Prev Med* 2002;23:51-61.

Rush AJ. Strategies and tactics in the management of maintenance treatment for depressed patients. *J Clin Psychiatry* 1999;60:21-26.

Rush AJ, Marangell LB, Sackeim HA, et al. Vagus nerve stimulation for treatment-resistant depression: a randomized, controlled acute phase trial. *Biol Psychiatry* 2005;58:347-54.

Rush AJ, Trivedi MH, Ibrahim HM, et al. The 16-item quick inventory of depressive symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): a psychometric evaluation in patients with chronic major depression. *Biol Psychiatry* 2003;54:573-83.

Rush AJ, Trivedi MH, Wisniewski SR, et al. Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report. *Am J Psychiatry* 2006;163:1905-17.

Rush AJ, Warden D, Wisniewski SR, et al. STAR*D: revising conventional wisdom. *CNS Drugs* 2009;23:627-47.

Sackeim HA. Magnetic stimulation therapy and ECT. *Convulsive Therapy* 1994;10:255-58.

Sackeim HA, Brannan SK, Rush AJ, et al. Durability of antidepressant response to vagus nerve stimulation (VNS™). *Int J Neuropsychopharmacol* 2007;10:817-26.

Sackheim HA, Haskett RF, Mulsant BH, et al. Continuation pharmacotherapy in the prevention of relapse following electroconvulsive therapy: a randomized controlled trial. *JAMA* 2001a;285:1299-1307.

Sackeim HA, Keilp JG, Rush AJ, et al. The effects of vagus nerve stimulation on cognitive performance in patients with treatment-resistant depression. *Neuropsychiatry Neuropsychol Behav Neurol* 2001b;14:53-62.

Sackeim HA, Rush AJ, George MS, et al. Vagus nerve stimulation (VNS™) for treatment-resistant depression: efficacy, side effects, and predictors of outcome. *Neuropsychopharmacology* 2001c;25:713-28.

Schatzberg AF, Rush AJ, Arnow BA, et al. Chronic depression: medication (nefazodone) or psychotherapy (CBASP) is effective when the other is not. *Arch Gen Psychiatry* 2005;62:513-20.

Schlaepfer TE, Frick C, Zobel A, et al. Vagus nerve stimulation for depression: efficacy and safety in a European study. *Psychol Med* 2008;38:651-61.

Schoenbaum M, Unützer J, McCaffrey D, et al. The effects of primary care depression treatment on patients' clinical status and employment. *Health Serv Res* 2002;37:1145-58.

Schoenbaum M, Unützer J, Sherbourne C, et al. Cost-effectiveness of practice-initiated quality improvement for depression: results of a randomized controlled trial. *JAMA* 2001;286:1325-30.

Schonfeld WH, Verboncoeur CJ, Fifer SK, et al. The functioning and well-being of patients with unrecognized anxiety disorders and major depressive disorder. *J Affect Disord* 1997;43:105-19.

Schramm E, Zobel I, Dykierek P, et al. Cognitive behavioral analysis system of psychotherapy versus interpersonal psychotherapy for early-onset chronic depression: a randomized pilot study. *J Affect Disord* 2011;129:109-16. www.icsi.org

References

- Schraufnagel TJ, Wagner AW, Miranda J, Roy-Byrne PP. Treating minority patients with depression and anxiety: what does the evidence tell us? *Gen Hosp Psychiatry* 2006;28:27-36.
- Schuch FB, Vasconcelos-Moreno MP, Borowsky C, Fleck MP. Exercise and severe depression: preliminary results of an add-on study. *J Affect Disord* 2011;133:615-18.
- Schulberg HC, Katon W, Simon GE, Rush AJ. Treating major depression in primary care practice: an update of the agency for health care policy and research practice guidelines. *Arch Gen Psychiatry* 1998;55:1121-27.
- Schwartz TL, Azhar N, Cole K, et al. An open-label study of adjunctive modafinil in patients with sedation related to serotonergic antidepressant therapy. *J Clin Psychiatry* 2004;65:1223-27.
- Segal ZV, Bieling P, Young T, et al. Antidepressant monotherapy vs sequential pharmacotherapy and mindfulness-based cognitive therapy, or placebo, for relapse prophylaxis in recurrent depression. *Arch Gen Psychiatry* 2010;67:1256-64.
- Shaffer JA, Edmondson D, Wasson LT, et al. Vitamin D supplementation for depressive symptoms: a systematic review and meta-analysis of randomized controlled trials. *Psychosom Med* 2014;76:190-96.
- Sharma V, Khan M, Smith A. A closer look at treatment resistant depression: is it due to a bipolar diathesis? *J Affective Disorders* 2005;84:251-57.
- Shelton RC, Osuntokun O, Heinloth AN, Corya SA. Therapeutic options for treatment-resistant depression. *CNS Drugs* 2010;24:131-61.
- Shinkai RS, Hatch JP, Schmidt CB, Sartori EA. Exposure to the oral side effects of medication in a community-based sample. *Spec Care Dentist* 2006;26:116-20.
- Silveira H, Moraes H, Oliveira N, et al. Physical exercise and clinically depressed patients: a systematic review and meta-analysis. *Neuropsychobiology* 2013;67:61-68.
- Simon GE, Katon WJ, VonKorff M, et al. Cost-effectiveness of a collaborative care program for primary care patients with persistent depression. *Am J Psychiatry* 2001a;158:1638-44.
- Simon GE, Manning WG, Katzelnick DJ, et al. Cost-effectiveness of systematic depression treatment for high utilizers of general medical care. *Arch Gen Psychiatry* 2001b;58:181-87.
- Simon GE, Savarino J. Suicide attempts among patients starting depression treatment with medications or psychotherapy. *Am J Psychiatry* 2007;164:1029-34.
- Simon GE, Van Korff M, Ludman EJ, et al. Cost-effectiveness of a program to prevent depression relapse in primary care. *Medical Care* 2002;40:941-50.
- Simon GE, Van Korff M, Rutter C, Wagner E. Randomised trial of monitoring, feedback, and management of care by telephone to improve treatment of depression in primary care. *BMJ* 2000;320:550-54.
- Siu AL, US Preventive Services Task Force (USPSTF), Bibbins-Domingo K, et al. Screening for depression in adults: U.S. preventive services task force recommendation statement. *JAMA* 2016;315:380-87.
- Sivojelezova A, Shuhaiber S, Sarkissian L, et al. Citalopram use in pregnancy: prospective comparative evaluation of pregnancy and fetal outcome. *Am J Obstet Gynecol* 2005;193:2004-09.
- Siwek M, Dudek D, Schlegel-Zawadzka M, et al. Serum zinc level in depressed patients during zinc supplementation of imipramine treatment. *J Affect Disord* 2010;126:447-52.
- Smith CA, Hay PP, Macpherson H. Acupuncture for depression. *Cochrane Database Syst Rev* 2010;(1):CD004046.
- Snieszek DP, Siddiqui IJ. Acupuncture for treating anxiety and depression in women: a clinical systematic

References

- review. *Med Acupunct* 2013;25:164-72.
- Solomon DA, Keller MB, Leon AC, et al. Multiple recurrences of major depressive disorder. *Am J Psychiatry* 2000;157:229-33.
- Sonawalla SB. Citalopram in the maintenance treatment of major depressive disorder. *J Clin Psychiatry* 2001;62:993.
- Sorrentino R. Performing capacity evaluations: what's expected from your consult: core components of a capacity evaluation are understanding, free choice, and reliability. *Curr Psychiatry* 2014;1:41-44.
- Spielmanns GI, Berman MI, Usitalo AN. Psychotherapy versus second-generation antidepressants in the treatment of depression: a meta-analysis. *J Nerv Ment Dis* 2011;199:142-49.
- Spier SP. Use of bupropion with SRIs and venlafaxine. *Depression and Anxiety* 1998;7:73-75.
- Spijker J, van Straten A, Bockting CL, et al. Psychotherapy, antidepressants, and their combination for chronic major depressive disorder: a systematic review. *Can J Psychiatry* 2013;58:386-92.
- Spitzer RL, Kroenke K, Williams JBW. Validation and utility of a self-report version of prime-md: the PHQ primary care study. *JAMA* 1999;282:1737-44.
- Steinert C, Hofmann M, Kruse J, Leichsenring F. Relapse rates after psychotherapy for depression – stable long-term effects? A meta-analysis. *J Affect Disord* 2014;168:107-18.
- Stewart DE. Depression during pregnancy. *N Engl J Med* 2011;365:1605-11.
- Stewart WF, Ricci JA, Chee E, et al. Cost of lost productive work time among US workers with depression. *JAMA* 2003;289:3135-44.
- Strine TW, Mokdad AH, Balluz LS, et al. Depression and anxiety in the United States: findings from the 2006 behavioral risk factor surveillance system. *Psychiatr Serv* 2008;59:1383-90.
- Szymkowicz SM, Finnegan N, Dale RM. A 12-month naturalistic observation of three patients receiving repeat intravenous ketamine infusions for their treatment-resistant depression. *J Affect Disord* 2013;147:416-20.
- Teasdale JD, Scott J, Moore RG, et al. How does cognitive therapy prevent relapse in residual depression: evidence from a controlled trial. *J Consult Clin Psychol* 2001;69:347-57.
- Thacher TD, Clarke BL. Vitamin D insufficiency. *Mayo Clin Proc* 2011;86:50-60.
- Thase ME, Blomgren SL, Birkett MA, et al. Fluoxetine treatment of patients with major depressive disorder who failed initial treatment with sertraline. *J Clin Psychiatry* 1997;58:16-21.
- Thase ME, Friedman ES, Biggs MM, et al. Cognitive therapy versus medication in augmentation and switch strategies as second-step treatments: a STAR*D report. *Am J Psychiatry* 2007;164:739-52.
- Thormahlen GM. Paroxetine using during pregnancy: is it safe? *Ann Pharmacother* 2006;40:1834-37.
- Trivedi MH. Tools and strategies for ongoing assessment of depression: a measurement-based approach to remission. *J Clin Psychiatry* 2009;70:26-31.
- Trivedi MH, Fava M, Wisniewski SR, et al. Medication augmentation after the failure of SSRIs for depression. *N Engl J Med* 2006a;354:1243-52.
- Trivedi MH, Kleiber BA. Algorithm for the treatment of chronic depression. *J Clin Psychiatry* 2001;62:22-29.
- Trivedi MH, Rush AJ, Wisniewski SR, et al. Evaluation of outcomes with citalopram for depression using

References

- measurement-based care in STAR*D: implications for clinical practice. *Am J Psychiatry* 2006b;163:28-40.
- Tschoppe P, Wolgin M, Pischon N, Kielbassa AM. Etiologic factors of hyposalivation and consequences for oral health. *Quintessence Int* 2010;41:321-33.
- UK ECT Review Group, The. Efficacy and safety of electroconvulsive therapy in depressive disorders: a systematic review and meta-analysis. *Lancet* 2003;361:799-808.
- U.S. Preventive Services Task Force. Screening and behavioral counseling interventions in primary care to reduce alcohol misuse: recommendation statement. *Am Fam Phys* 2014;89.
- Unützer J, Katon W, Callahan CM, et al. Collaborative care management of late-life depression in the primary care setting: a randomized controlled trial. *JAMA* 2002;288:2836-45.
- Unützer J, Katon WJ, Fan MY, et al. Long-term cost effects of collaborative care for late-life depression. *Am J Manag Care* 2008;14:95-100.
- Unützer J, Tang L, Oishi S, et al. Reducing suicidal ideation in depressed older primary care patients. *J Am Geriatr Soc* 2006;54:1550-56.
- van Dijk SE, Pols AD, Adriaanse MC, et al. Cost-effectiveness of a stepped-care intervention to prevent major depression in patients with type 2 diabetes mellitus and/or coronary heart disease and subthreshold depression: design of a cluster-randomized controlled trial. *BMC Psychiatry* 2013;13:128.
- van Hees ML, Rotter T, Ellermann T, Evers SM. The effectiveness of individual interpersonal psychotherapy as a treatment for major depressive disorder in adult outpatients: a systematic review. *BMC Psychiatry* 2013;13:22.
- Vanelli M, Coca-Perraillon M. Role of patient experience in antidepressant adherence: a retrospective data analysis. *Clin Ther* 2008;30:1737-45.
- Vesga-López O, Blanco C, Keyes K, et al. Psychiatric disorders in pregnant and postpartum women in the United States. *Arch Gen Psychiatry* 2008;65:805-15.
- Vittengl JR, Clark LA, Jarrett RB. Continuation-phase cognitive therapy's effects on remission and recovery from depression. *J Consult Clin Psychol* 2009;77:367-71.
- von Wolff A, Hölzel LP, Westphal A, et al. Selective serotonin reuptake inhibitors and tricyclic antidepressants in the acute treatment of chronic depression and dysthymia: a systematic review and meta-analysis. *J Affect Disord* 2013;144:7-15.
- Wang F, Lee EO, Wu T, et al. The effects of tai chi on depression, anxiety, and psychological well-being: a systematic review and meta-analysis. *Int J Behav Med* 2014;21:605-17.
- Wang PS, Simon GE, Avorn J, et al. Telephone screening, outreach, and care management for depressed workers and impact on clinical and work productivity outcomes: a randomized controlled trial. *JAMA* 2007;298:1401-11.
- Watkins S, Meltzer-Brody S, Zolnoun D, Stuebe A. Early breastfeeding experiences and postpartum depression. *Obstet Gynecol* 2011;118:214-21.
- Watson LC, Zimmerman S, Cohen LW, Dominik R. Practical depression screening in residential care/assisted living: five methods compared with gold standard diagnoses. *Am J Geriatr Psychiatry* 2009;17:556-64.
- Wenzel RG, Tepper S, Korab WE, Freitag F. Serotonin syndrome risks when combining SSRI/SNRI drugs and triptans: is the FDA's alert warranted? *Ann Pharmacother* 2008;42:1692-96.
- Wernicke JF, Pritchett YL, D'Souza DN, et al. A randomized controlled trial of duloxetine in diabetic

References

- peripheral neuropathic pain. *Neurology* 2006;67:1411-20.
- Whiting PF, Wolff RF, Deshpande S, et al. Cannabinoids for medical use: a systematic review and meta-analysis. *JAMA* 2015;313:2456-73.
- Whooley MA, de Jonge P, Vittinghoff E, et al. Depressive symptoms, health behaviors, and risk of cardiovascular events in patients with coronary heart disease. *JAMA* 2008;300:2379-88.
- Whooley MA, Simon GE. Managing depression in medical outpatients. *N Engl J Med* 2000;343:1942-50.
- Wiersma JE, Van Schaik DJ, Hoogendorn AW, et al. The effectiveness of the cognitive behavioral analysis system of psychotherapy for chronic depression: a randomized controlled trial. *Psychother Psychosom* 2014;83:263-69.
- Williams Jr JW, Noel PH, Cordes JA, et al. Is this patient clinically depressed? *JAMA* 2002;287:1160-70.
- Wisner KL, Sit DKY, Hanusa BH, et al. Major depression and antidepressant treatment: impact on pregnancy and neonatal outcomes. *Am J Psychiatry* 2009;166:557-66.
- Wittkamp KA, Naeije L, Schene AH, et al. Diagnostic accuracy of the mood module of the patient health questionnaire: a systematic review. *Gen Hosp Psychiatry* 2007;29:388-95.
- Wright BM, Eiland III EH, Lorenz R. Augmentation with atypical antipsychotics for depression: a review of evidence-based support from the medical literature. *Pharmacotherapy* 2013;33:344-59.
- Wulsin LR, Singal BM. Do depressive symptoms increase the risk for the onset of coronary disease? A systematic quantitative review. *Psychosom Med* 2003;65:201-10.
- Wulsin L, Somoza E, Heck J. The feasibility of using the Spanish PHQ-9 to screen for depression in primary care in Honduras. *Prim Care Companion J Clin Psychiatry* 2002;4:191-95.
- Yates WR, Mitchell J, Rush AJ, et al. Clinical features of depression in outpatients with and without co-occurring general medical conditions in STAR*D: confirmatory analysis. *Prim Care Companion J Clin Psychiatry* 2007;9:7-15.
- Yeung A, Fung F, Yu SC, et al. Validation of the patient health questionnaire-9 for depression screening among Chinese Americans. *Compr Psychiatry* 2008;49:211-17.
- Yonkers KA, Wisner KL, Stewart DE, et al. The management of depression during pregnancy: a report from the American psychiatric association and the American college of obstetricians and gynecologists. *Gen Hosp Psychiatry* 2009;31:403-13.
- Zhornitsky S, Potvin S, Motesafi H, et al. Dose-response and comparative efficacy and tolerability of quetiapine across psychiatric disorders: a systematic review of the placebo-controlled monotherapy and add-on trials. *Int Clin Psychopharmacol* 2011;26:183-92.
- Zivin K, Pfeiffer PN, Bohnert AS, et al. Evaluation of the FDA warning against prescribing citalopram at doses exceeding 40 mg. *Am J Psychiatry* 2013;170:642-50.