

The subdivisions of this section are:

- References

## References

- Altamimi S, Robertson G, Jastaniah W, et al. Single-dose oral dexamethasone in the emergency management of children with exacerbations of mild to moderate asthma. *Pediatr Emerg Care* 2006;22:786-93.
- Cowan DC, Hewitt RS, Cowan JO, et al. Exercise-induced wheeze: fraction of exhaled nitric oxide-directed management. *Respirology* 2010;15:683-90.
- Cronin JJ, McCoy S, Kennedy U, et al. A randomized trial of single-dose oral dexamethasone versus multidose prednisolone for acute exacerbations of asthma in children who attend the emergency department. *Ann Emerg Med* 2016;67:593-601.
- Cross KP, Paul RI, Goldman RD. Single-dose dexamethasone for mild-to-moderate asthma exacerbations: effective, easy, and acceptable. *Can Fam Physician* 2011;57:1134-36.
- de Jongste JC, Carraro S, Hop WC, et al. Daily telemonitoring of exhaled nitric oxide and symptoms in the treatment of childhood asthma. *Am J Respir Crit Care Med* 2009;179:93-97.
- Dinh-Xuan AT, Annesi-Maesano I, Berger P, et al. Contribution of exhaled nitric oxide measurement in airway inflammation assessment in asthma. A position paper from the French speaking respiratory society. *Rev Mal Respir* 2015;32:193-215.
- Dweik RA, Boggs PB, Erzurum SC, et al. An official ATS clinical practice guideline: interpretation of exhaled nitric oxide levels (FE<sub>NO</sub>) for clinical applications. *Am J Respir Crit Care Med* 2011;184:602-15.
- Feitosa LA, Dornelas de Andrade A, Reinaux CM, Britto MC. Diagnostic accuracy of exhaled nitric oxide in exercise-induced bronchospasm: systematic review. *Rev Port Pneumol* 2012;18:198-204.
- Food and Drug Administration. FDA announces new safety controls for long-acting beta agonists, medications used to treat asthma. 2010.
- Gomersal T, Harnan S, Essat M, et al. A systematic review of fractional exhaled nitric oxide in the routine management of childhood asthma. *Pediatr Pulmonol* 2016;51:316-28.
- Gruchalla RS, Sampson HA, Matsui E, et al. Asthma morbidity among inner-city adolescents receiving guidelines-based therapy: role of predictors in the setting of high adherence. *J Allergy Clin Immunol* 2009;124:213-21.
- Guo Z, Wang Y, Xing G, Wang X. Diagnostic accuracy of fractional exhaled nitric oxide in asthma: a systematic review and meta-analysis of prospective studies. *J Asthma* 2016;53:404-12.
- Harnan S, Essat M, Gomersall T, et al. Exhaled nitric oxide for the diagnosis of asthma in adults and children: a systematic review. *Value Health* 2015a;18:A345.
- Harnan SE, Tappenden P, Essat M, et al. Measurement of exhaled nitric oxide concentration in asthma: a systematic review and economic evaluation of NIOX MINO, NIOX VERO and NObreath. *Health Technol Assess* 2015b;19:1-330.
- Jartti T, Wendelin-Saarenhovi M, Heinonen I, et al. Childhood asthma management guided by repeated FeNO measurements: a meta-analysis. *Paediatr Respir Rev* 2012;13:178-83.
- Keeney GE, Gray MP, Morrison AK, et al. Dexamethasone for acute asthma exacerbations in children: a meta-analysis. *Pediatrics* 2014;133:493-99.
- Korevaar DA, Westerhof GA, Wang J, et al. Diagnostic accuracy of minimally invasive markers for detection of airway eosinophilia in asthma: a systematic review and meta-analysis. *Lancet Respir Med* 2015;3:290-300.

**References**

- Kravitz J, Dominici P, Ufberg J, et al. Two days of dexamethasone versus 5 days of prednisone in the treatment of acute asthma: a randomized controlled trial. *Ann Emerg Med* 2011;58:200-04.
- LaForce C, Brooks E, Herje N, et al. Impact of exhaled nitric oxide measurements on treatment decisions in an asthma specialty clinic. *Ann Allergy Asthma Immunol* 2014;113:619-23.
- Lehtimäki L, Csonka P, Mäkinen E, et al. Predictive value of exhaled nitric oxide in the management of asthma: a systematic review. *Eur Respir J* 2016;48:706-14.
- Lemiere C, Nguyen S, Sava F, et al. Occupational asthma phenotypes identified by increased fractional exhaled nitric oxide after exposure to causal agents. *J Allergy Clin Immunol* 2014;134:1063-67.
- Lu M, Wu B, Che D, et al. FeNO and asthma treatment in children: a systematic review and meta-analysis. *Medicine (Baltimore)* 2015;94:e347.
- Meyer JS, Riese J, Biondi E. Is dexamethasone an effective alternative to oral prednisone in the treatment of pediatric asthma exacerbations? *Hosp Pediatr* 2014;4:172-80.
- Peirsman EJ, Carvelli TJ, Hage PY, et al. Exhaled nitric oxide in childhood allergic asthma management: a randomised controlled trial. *Pediatr Pulmonol* 2014;49:624-31.
- Petsky HL, Cates CJ, Li A, et al. Tailored interventions based on exhaled nitric oxide versus clinical symptoms for asthma in children and adults. *Cochrane Database Syst Rev* 2009;(4):CD006340.
- Scott HA, Latham JR, Callister R, et al. Acute exercise is associated with reduced exhaled nitric oxide in physically inactive adults with asthma. *Ann Allergy Asthma Immunol* 2015;114:470-79.
- Syk J, Malinowski A, Johansson G, et al. Anti-inflammatory treatment of atopic asthma guided by exhaled nitric oxide: a randomized, controlled trial. *J Allergy Clin Immunol Pract* 2013;1:639-48.
- Tang S, Xie Y, Yuan C, et al. Fractional exhaled nitric oxide for the diagnosis of childhood asthma: a systematic review and meta-analysis. *Clin Rev Allergy Immunol* 2016.
- Voorend-van Bergen S, Vaessen-Verberne AA, Landstra AM, et al. Fractional exhaled nitric oxide monitoring does not improve asthma management in children with concordant and discordant asthma phenotypes. *Am J Respir Crit Care Med* 2015;192:1016-18.