

The subdivisions of this section are:

- References

## References

Links are provided for those new references added to this edition (author name is highlighted in blue).

- [Acharya SD, Elci OU, Sereika SM, et al.](#) Using a personal digital assistant for self-monitoring influences diet Sixth Edition/January 2016 in comparison to a standard paper record among overweight/obese adults. *J Am Diet Assoc* 2011;111:583-88.
- [Allen JG, Flanigan SS, LeBlanc M, et al.](#) Flavoring chemicals in e-cigarettes: diacetyl, 2,3-pentanedione, and acetoin in a sample of 51 products, including fruit-, candy-, and cocktail-flavored e-cigarettes. *Environ Health Perspect* 2015.
- [Alvarez-Bueno C, Rodriguez-Martin B, Garcia-Ortiz L, et al.](#) Effectiveness of brief interventions in primary health care settings to decrease alcohol consumption by adult non-dependent drinkers: a systematic review of systematic reviews. *Prev Med* 2015;76 Suppl:S33-38.
- [Altman NG, Schopfer E, Jackson N, et al.](#) Sleep duration versus sleep insufficiency as predictors of cardiometabolic health outcomes. *Sleep Med* 2012 ;13:1261-70.
- [American Psychiatric Association.](#) *In Diagnostic and Statistical Manual of Mental Disorders DSM-5.* Fifth Edition. Washington, DC/London, England. 2013.
- Ammerman AS, Lindquist CH, Lohr KN, Hersey J. The efficacy of behavioral interventions to modify dietary fat and fruit and vegetable intake: a review of the evidence. *Prev Med* 2002;35:25-41.
- Armitage CJ. Evidence that implementation intentions reduce dietary fat intake: a randomized trial. *Health Psychol* 2004;23:319-23.
- [Ashford S, Edmunds J, French DP.](#) What is the best way to change self-efficacy to promote lifestyle and recreational physical activity? A systematic review with meta-analysis. *Br J Health Psychol* 2010;15:265-88.
- [Babor TF, Higgins-Biddle JC.](#) *In Brief Intervention for Hazardous and Harmful Drinking: A Manual for Use in Primary Care.* World Health Organization: The Department of Mental Health and Substance Dependence. 2001a.
- [Babor TF, Higgins-Biddle JC, Saunders JB, Monteiro MG.](#) AUDIT, the alcohol use disorders test: guidelines for use in primary care. World Health Organization: The Department of Mental Health and Substance Dependence. 2001b.
- Babor TF, McRee BG, Kassebaum PA, et al. Screening, brief intervention, and referral to treatment (SBIRT): toward a public health approach to the management of substance abuse. *Subst Abuse* 2007;28:7-30.
- [Bellavia A, Larsson SC, Bottai M, et al.](#) Fruit and vegetable consumption and all-cause mortality: a dose-response analysis. *Am J Clin Nutr* 2013;98:454-59.
- Bertholet N, Daepfen JB, Wietlisbach V, et al. Reduction by alcohol consumption by brief alcohol intervention in primary care: systematic review and meta-analysis. *Arch Intern Med* 2005;165:986-95.
- [Bhattarai N, Prevost AT, Wright AJ, et al.](#) Effectiveness of interventions to promote healthy diet in primary care: a systematic review and meta-analysis of randomized controlled trials. *BMC Public Health* 2013;13:1203.
- [Bolier L, Haverman M, Westerhof GJ, et al.](#) Positive psychology interventions: a meta-analysis of randomized controlled studies. *BMC Public Health* 2013;13:119.
- Boutelle KN, Kirschenbaum DS. Further support for consistent self-monitoring as a vital component of successful weight control. *Obes Res* 1998;6:219-24.

## References

- Bovend'Eerdt TJ, Botell RE, Wade DT. Writing SMART rehabilitation goals and achieving goal attainment scaling: a practical guide. *Clin Rehabil* 2009;23:352-61.
- Boyle R, Solberg L, Fiore M. Use of electronic health records to support smoking cessation. *Cochrane Database Syst Rev* 2014;(12):CD008743.
- Bradley KA, Bush KR, Epler AJ, et al. Two brief alcohol-screening tests From the Alcohol Use Disorders Identification Test (AUDIT): validation in a female Veterans Affairs patient population. *Arch Intern Med* 2003;163:821-29.
- Bravata DM, Smith-Spangler C, Sundaram V, et al. Using pedometers to increase physical activity and improve health: a systematic review. *JAMA* 2007;298:2296-2304.
- Brownell KD, Farley T, Willett WC, et al. The public health and economic benefits of taxing sugar-sweetened beverages. *N Engl J Med* 2009;361:1599-1605.
- Bully P, Sanchez A, Zabaleta-del-Olmo E, et al. Evidence from interventions based on theoretical models for lifestyle modification (physical activity, diet, alcohol and tobacco use) in primary care settings: a systematic review. *Prev Med* 2015;76:S76-S93.
- Burke LE, Conroy MB, Sereika SM, et al. The effect of electronic self-monitoring on weight loss and dietary intake: a randomized behavioral weight loss trial. *Obesity* 2011;19:338-44.
- Bush K, Kivlahan DR, McDonell MB, et al. The AUDIT alcohol consumption questions (AUDIT-C). *Arch Intern Med* 1998;158:1789-95.
- Butler CC, Rollnick S, Cohen D, et al. Motivational advice vs brief consulting for smokers in general practice: a randomised trial. *Brit J Gen Pract* 1999;49:611-16.
- Canagasaby A, Vinson DC. Screening for hazardous or harmful drinking using one or two quantity-frequency questions. *Alcohol Alcohol* 2005;40:208-13.
- Centers for Disease Control and Prevention. Effects of short sleep duration on daily activities – United States, 2005-2008. *Morbidity and Mortality Weekly Report* 2011a;60:239-42.
- Centers for Disease Control and Prevention. Smoking and Overall Health. From the Surgeon General's Report. Fact Sheet for Public Health Officials. Available at: [http://www.cdc.gov/tobacco/data\\_statistics/sgr/50th-anniversary/pdfs/fs\\_smoking\\_overall\\_health\\_508.pdf](http://www.cdc.gov/tobacco/data_statistics/sgr/50th-anniversary/pdfs/fs_smoking_overall_health_508.pdf). 2014.
- Centers for Disease Control and Prevention. Unhealthy sleep-related behaviors – 12 states 2009. *Morbidity and Mortality Weekly Report* 2011b;60:234-38.
- Chapman CD, Benedict C, Brooks SJ, Schioth HB. Lifestyle determinants of the drive to eat: a meta-analysis. *Am J Clin Nutr* 2012;96:492-97.
- Chen L, Appel LJ, Loria C, et al. Reduction in consumption of sugar-sweetened beverages is associated with weight loss: the PREMIER trial. *Am J Clin Nutr* 2009;89:1299-306.
- Chen L, Caballero B, Mitchell DC, et al. Reducing consumption of sugar-sweetened beverages is associated with reduced blood pressure: a prospective study among United States adults. *Circulation* 2010;121:2398-2406.
- Chung F, Yegneswaran B, Liao P, et al. STOP questionnaire: a tool to screen patients for obstructive sleep apnea. *Anesthesiology* 2008;108:812-21.
- Daviglus ML, Stamler J, Pirzada A, et al. Favorable cardiovascular risk profile in young women and long-term risk of cardiovascular and all-cause mortality. *JAMA* 2004;292:1588-92.
- Dietary Guidelines Advisory Committee. Scientific report of the 2015 dietary guidelines advisory committee. Available at: <http://health.gov/dietaryguidelines/2015-scientific-report/PDFs/Scientific-Report-of-the-2015-Dietary-Guidelines-Advisory-Committee.pdf>. 2015.

## References

- Donoghue K, Patton R, Phillips T, et al. The effectiveness of electronic screening and brief intervention for reducing levels of alcohol consumption: a systematic review and meta-analysis. *J Med Internet Res* 2014;16:e142.
- Dunn C, Deroo L, Rivara F. The use of brief interventions adapted from motivational interviewing across behavioral domains: a systematic review. *Addiction* 2001;96:1725-42.
- Eakin EG, Lawler SP, Vandelanotte C, Owen N. Telephone interventions for physical activity and dietary behavior change: a systematic review. *Am J Prev Med* 2007;32:419-34.
- Ebbert JO, Agunwamba AA, Rutten LJ. Counseling patients on the use of electronic cigarettes. *Mayo Clin Proc* 2015;90:128-34.
- Emmons RA, McCullough ME. Counting blessings versus burdens: an experimental investigation of gratitude and subjective well-being in daily life. *J Pers Soc Psychol* 2003;84:377-89.
- Estruch R, Ros E, Salas-Salvadó J, et al. Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med* 2013;368:1279-89.
- Feart C, Samieri C, Rondeau V, et al. Adherence to a Mediterranean diet, cognitive decline, and risk of dementia. *JAMA* 2009;302:638-48.
- Frank D, DeBenedetti AF, Volk RJ, et al. Effectiveness of the AUDIT-C as a screening tool for alcohol misuse in three race/ethnic groups. *J Gen Intern Med* 2008;23:781-87.
- Fredrickson BL. The role of positive emotions in positive psychology: the broaden-and-build theory of positive emotions. *Am Psychol* 2001;56:218-26.
- Fredrickson BL, Cohn MA, Coffey KA, et al. Open hearts build lives: positive emotions, induced through loving-kindness meditation, build consequential personal resources. *J Pers Soc Psychol* 2008;95:1045-62.
- Fries E, Edinboro P, McClish D, et al. Randomized trial of a low-intensity dietary intervention in rural residents: the rural clinician cancer prevention project. *Am J Prev Med* 2005;28:162-68.
- Fung TT, Chiuve SE, McCullough ML, et al. Adherence to a DASH-style diet and risk of coronary heart disease and stroke in women. *Arch Intern Med* 2008;168:713-20.
- Gable SL, Reis HT, Impett EA, Asher ER. What do you do when things go right? The intrapersonal and interpersonal benefits of sharing positive events. *J Pers Soc Psychol* 2004;87:228-45.
- Gander F, Proyer RT, Ruch W, Wyss T. Strength-based positive interventions: further evidence for their potential in enhancing well-being and alleviating depression. *J Happiness Stud* 2013;14:1241-59.
- Gans KM, Risica PM, Wylie-Rosett J, et al. Development and evaluation of the nutrition component of the rapid eating and activity assessment for patients (REAP): a new tool for primary care providers. *J Nutr Educ Behav* 2006;38:286-92.
- Gans KM, Ross E, Barner CW, et al. REAP and WAVE: new tools to rapidly assess/discuss nutrition with patients. *J Nutr* 2003;133:556S-62S.
- Glasgow RE, Davis CL, Funnell MM, Beck A. Implementing practical interventions to support chronic illness self-management in health care settings: lessons learned and recommendations. *Jt Comm J Qual Saf* 2003;29:563-74.
- Goetzel RZ, Ozminkowski RJ, Bruno JA, et al. The long-term impact of Johnson & Johnson's health & wellness program on employee health risks. *J Occup Environ Med* 2002;44:417-24.
- Gold D, Grossmeier J. Using an HRA as an evaluation tool. *Wellness Management* 2005;20:1-5.

## References

- Gomel M, Oldenburg B, Simpson JM, Owen N. Work-site cardiovascular risk reduction: a randomized trial of health risk assessment, education, counseling, and incentives. *Am J Public Health* 1993;83:1231-38.
- Grigsby-Toussaint DS, Turi KN, Krupa M, et al. Sleep insufficiency and the natural environment: results from the US behavioral risk factor surveillance system survey. *Prev Med* 2015;78:78-84.
- Harmon BE, Boushey CJ, Shvetsov YB, et al. Associations of key diet-quality indexes with mortality in multiethnic cohort: the dietary patterns methods project. *Am J Clin Nutr* 2015;101:587-97.
- Hedrick VE, Savla J, Comber DL, et al. Development of a brief questionnaire to assess habitual beverage intake (BEVQ-15): sugar sweetened beverages and total beverage energy intake. *J Acad Nutr Diet* 2012;112:840-49.
- Hirshkowitz M, Whiton K, Albert S, et al. The national sleep foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health* 2015;1:40-43.
- Institute of Medicine (US) Committee on Sleep Medicine and Research. Sleep disorders and sleep deprivation: an unmet public health problem. Washington, DC: National Academies Press (US). 2006.
- Jenkins DJA, Kendall CWC, Faulkner DA, et al. Assessment of the longer-term effects of a dietary portfolio of cholesterol-lowering foods in hypercholesterolemia. *Am J Clin Nutr* 2006;83:582-91.
- Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep* 1991;14:540-45.
- Johnson M, Jackson R, Guillaume L, et al. Barriers and facilitators to implementing screening and brief intervention for alcohol misuse: a systematic review of qualitative evidence. *J Public Health* 2010;33:412-21.
- Jonas DE, Garbutt JC, Brown JM, et al. Screening, behavioral counseling, and referral in primary care to reduce alcohol misuse. Rockville (MD): Agency for Health care Research and Quality (US). 2012.
- Kang M, Marshall SJ, Barreira TV, Lee JO. Effect of pedometer-based physical activity interventions: a meta-analysis. *Res Q Exerc Sport* 2009;80: 648-55.
- Kaner EF, Dickinson HO, Beyer F, et al. The effectiveness of brief alcohol interventions in primary care settings: a systematic review. *Drug Alcohol Rev* 2009;28:301-23.
- Kant AK, Schatzkin A, Graubard BI, Schairer C. A prospective study of diet quality and mortality in women. *JAMA* 2000;283:2109-15.
- Kastorini CM, Milionis HJ, Esposito K, et al. The effect of Mediterranean diet on metabolic syndrome and its components: a meta-analysis of 50 studies and 534,906 individuals. *J Am Coll Cardiol* 2011;57:1299-313.
- Keyes CL, Simoes EJ. To flourish or not: positive mental health and all-cause mortality. *Am J Public Health* 2012;102:2164-72.
- Kim JJ, Yadav D, Ahn SV, et al. A prospective study of total sleep duration and incident metabolic syndrome: the ARIRANG study. *Sleep Med* 2015;16:1511-15.
- Knutson KL. Sleep duration and cardiometabolic risk: a review of the epidemiologic evidence. *Best Pract Res Clin Endocrinol Metab* 2010; 24:731-43.
- Koloverou E, Panagiotakos DB, Pitsavos C, et al. Adherence to Mediterranean diet and 10-year incidence (2002-2012) of diabetes: correlations with inflammatory and oxidative stress biomarkers in the ATTICA cohort study. *Diabetes Metab Res Rev* 2016;32:73-81.
- Kottke TE, Solberg LI, Brekke ML, et al. Will patient satisfaction set the preventive services implementation agenda? *Am J Prev Med* 1997;13:309-16.

## References

- Krueger PM, Friedman EM. Sleep duration in the United States: a cross-sectional population-based study. *Am J Epidemiol* 2009;169:1052-63.
- Lauer MS. Primary prevention of atherosclerotic cardiovascular disease: the high public burden of low individual risk. *JAMA* 2007;297:1376-78.
- Lavizzo-Mourey R. Childhood obesity: what it means for clinicians. *JAMA* 2007;298:920-22.
- LeFevre ML. Behavioral counseling to promote a healthful diet and physical activity for cardiovascular disease prevention in adults with cardiovascular risk factors: U.S. preventive services task force recommendation statement. *Ann Intern Med* 2014;161:587-93.
- Leventhal AM, Strong DR, Kirkpatrick MG, et al. Association of electronic cigarette use with initiation of combustible tobacco product smoking in early adolescence. *JAMA* 2015;314:700-07.
- Libotte E, Siegrist M, Bucher T. The influence of plate size on meal consumption.: literature review and experiment. *Appetite* 2014;82:91-96.
- Lieffers JR, Hanning RM. Dietary assessment and self-monitoring with nutrition applications for mobile devices. *Can J Diet Pract Res* 2012;73:e253-60.
- Lindson-Hawley N, Thompson TP, Begh R. Motivational interviewing for smoking cessation. *Cochrane Database Syst Rev* 2015;(3):CD006936.
- Lindström J, Louheranta A, Mannelin M, et al. The Finnish diabetes prevention study (DPS): lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care* 2003;26:3230-36. (High Quality Evidence)
- Lyubomirsky S, Layous K. How do simple positive activities increase well-being? *Current Directions in Psychological Science* 2013;22:57-62.
- Lyzwinski LN. A systematic review and meta-analysis of mobile devices and weight loss with an intervention content analysis. *J Pers Med* 2014;4:311-85.
- Maderuelo-Fernandez JA, Recio-Rodriguez JI, Patino-Alonso MC, et al. Effectiveness of interventions applicable to primary health care settings to promote Mediterranean diet or healthy eating adherence in adults: a systematic review. *Prev Med* 2015;76:S39-55.
- Madras BK, Compton WM, Avula D, et al. Screening, brief interventions, referral to treatment (SBIRT) for illicit drug and alcohol use at multiple health care sites: comparison at intake and 6 months later. *Drug Alcohol Depend* 2009;99:280-95.
- Maes S, Verhoeven C, Kittel F, Scholten H. Effects of a Dutch work-site wellness-health program: the Brabantia project. *Am J Public Health* 1998;88:1037-41.
- Maisto SA, Saitz R. Alcohol use disorders: screening and diagnosis. *Am J Addict* 2003;12:S12-S25.
- Malik VS, Popkin BM, Bray GA, et al. Sugar-sweetened beverages and risk of metabolic syndrome and type 2 diabetes: a meta-analysis. *Diabetes Care* 2010;33:2477-83.
- Malik VS, Schulze MB, Hu FB. Intake of sugar-sweetened beverages and weight gain: a systematic review. *Am J Clin Nutr* 2006;84:274-88.
- March S, Torres E, Ramos M, et al. Adult community health-promoting interventions in primary health care: a systematic review. *Prev Med* 2015;76:S94-S104.
- McGinnis JM, Foege WH. Actual causes of death in the United States. *JAMA* 1993;270:2207-12.
- McRobbie H, Bullen C, Hartmann-Boyce J, Hajek P. Electronic cigarettes for smoking and reduction. *Cochrane Database Syst Rev* 2014;(12):CD010216.



## References

- Moyer VA. Behavioral counseling interventions to promote a healthful diet and physical activity for cardiovascular disease prevention in adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2012;157:367-72.
- Moyer VA. Screening and behavioral counseling interventions in primary care to reduce alcohol misuse: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2013;159:210-18.
- National Institute for Health and Care Excellence (NICE). Physical activity: brief advice for adults in primary care. 2013.
- National Institute on Alcohol Abuse and Alcoholism. Helping patients who drink too much: a clinician's guide. 2005.
- National Prevention Council. National preventive strategy: active living. Available at: <http://www.surgeongeneral.gov/priorities/prevention/strategy/active-living.pdf>. 2014.
- Netzer NC, Stoohs RA, Netzer CM, et al. Using the Berlin questionnaire to identify patients at risk for the sleep apnea syndrome. *Ann Intern Med* 1999;131:485-91.
- Nothwehr F, Snetselaar L, Yang J, Wu H. Stage of change for healthful eating and use of behavioral strategies. *J Am Diet Assoc* 2006;106:1035-41.
- Obarzanek E, Sacks FM, Vollmer WM, et al. Effects on blood lipids of a blood pressure-lowering diet: the dietary approaches to stop hypertension (DASH) trial. *Am J Clin Nutr* 2001;74:80-89.
- O'Connor AM, Stacey D, Barry MJ, et al. Do patient decision aids meet effectiveness criteria of the international patient decision aid standards collaboration? A systematic review and meta-analysis. *Med Decis Making* 2007;27:554-74.
- Orr KK, Asal NJ. Efficacy of electronic cigarettes for smoking cessation. *Ann Pharmacother* 2014;48:1502-06.
- Ouweneel E, Le Blanc PM, Schaufeli WB. On being grateful and kind: results of two randomized controlled trials on study-related emotions and academic engagement. *J Psychol* 2014;148:37-60.
- Oyebode O, Gordon-Dseagu V, Walker A, Mindell JS. Fruit and vegetable consumption and all-cause cancer and CVD mortality: analysis of health survey for England data. *J Epidemiol Community Health* 2014;68:856-62.
- Panagiotakos DB, Pitsavos C, Skoumas Y, Stefanadis C. The association between food patterns and the metabolic syndrome using principal components analysis: the ATTICA Study. *J Am Diet Assoc* 2007;107:979-87.
- Patnode CD, Henderson JT, Thompson JH, et al. Behavioral counseling and pharmacotherapy interventions for tobacco cessation in adults, including pregnant women: a review for the U.S. Preventive Services Task Force. Rockville, MD: Agency for Health care Research and Quality. 2015.
- Paxton AE, Strycker LA, Toobert DJ, et al. Starting the conversation: performance of a brief dietary assessment and intervention tool for health professionals. *Am J Prev Med* 2011;40:67-71.
- Pignone MP, Ammerman A, Fernandez L, et al. Counseling to promote a healthy diet in adults: a summary of the evidence for the U.S. preventive services task force. *Am J Prev Med* 2003;24:75-92.
- Pollak KI, Alexander SC, Coffman CJ, et al. Physician communication techniques and weight loss in adults: project CHAT. *Am J Prev Med* 2010;39:321-28.
- Pomerleau J, Lock K, Knai C, McKee M. Interventions designed to increase adult fruit and vegetable intake can be effective: a systematic review of the literature. *J Nutr* 2005;135:2486-95.

## References

- Prochaska JO, Velicer WF, Redding C, et al. Stage-based expert systems to guide a population of primary care patients to quit smoking, eat healthier, prevent skin cancer, and receive regular mammograms. *Prev Med* 2005;41:406-16.
- Resnicow K, Dilorio C, Soet JE, et al. Motivational interviewing in health promotion: it sounds like something is changing. *Health Psych* 2002;21:444-51.
- Resnicow K, McCarty F, Baranowski T. Are precontemplators less likely to change their dietary behavior? A prospective analysis. *Health Educ Res* 2003;18:693-705.
- Richmond RL, Kehoe L, Hailstone S, et al. Quantitative and qualitative evaluations of brief interventions to change excessive drinking, smoking and stress in the police force. *Addiction* 1999;94:1509-21.
- Riemsma RP, Pattenden J, Bridle C, et al. Systematic review of the effectiveness of stage based interventions to promote smoking cessation. *BMJ* 2003;326:1175.
- [Robinson E, Nolan S, Tudur-Smith C, et al.](#) Will smaller plates lead to smaller waists? A systematic review and meta-analysis of the effect that experimental manipulation of dishware size has on energy consumption. *Obes Rev* 2014;15:812-21.
- Rubak S, Sandboek A, Lauritzen T, Christensen B. Motivational interviewing: a systematic review and meta-analysis. *Brit J Gen Pract* 2005;55:513:305-12.
- [Sanchez A, Bully P, Martinez C, Grandes G.](#) Effectiveness of physical activity promotion interventions in primary care: a review of reviews. *Prev Med* 2014;76:S56-S67.
- [Schulte B, O'Donnell AJ, Kastner S, et al.](#) Alcohol screening and brief intervention in workplace settings and social services: a comparison of literature. *Front Psychiatry* 2014;5:1-9.
- [Senthilvel E, Auckley D, Dasarathy J.](#) Evaluation of sleep disorder in primary care setting: history taking compared to questionnaires. *J Clin Sleep Med* 2011;7:41-48.
- [Shankar A, Syamala S, Kalidindi S.](#) Insufficient rest or sleep and its relation to cardiovascular disease, diabetes and obesity in a national, multiethnic sample. *PLoS ONE* 2010;5:e14189.
- [Shrestha N, Ijaz S, Kukkonen-Harjula KT, et al.](#) Workplace interventions for reducing sitting at work. *Cochrane Database Syst Rev* 2015;1:CD010912.
- [Sin NL, Lyubomirsky S.](#) Enhancing well-being and alleviating depressive symptoms with positive psychology interventions: a practice-friendly meta-analysis. *J Clin Psychol* 2009;65:467-87.
- [Smoking, nutrition, alcohol, physical activity \(SNAP\): a population health guide to behavioural risk factors in general practice.](#) 2nd Edition. Melbourne: The Royal Australian College of General Practitioners. 2015.
- [Siu AL.](#) Behavioral and pharmacotherapy interventions for tobacco smoking cessation in adults, including pregnant women: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 2015;163:622-34.
- [Smith PC, Schmidt SM, Allensworth-Davides D, Saitz R.](#) Primary care validation of a single-question alcohol screening test. *J Gen Intern Med* 2009;24:783-88.
- [Sofi F, Abbate R, Gensini GF, Casini A.](#) Accruing evidence on benefits of adherence to the Mediterranean diet on health: an updated systematic review and meta-analysis. *Am J Clin Nutr* 2010;92:1189-96.
- Sorensen G, Stoddard A, Hunt MK, et al. The effects of a health promotion – health protection intervention on behavior change: the WellWorks study. *Am J Public Health* 1998;88:1685-90.
- Sorensen G, Thompson B, Glanz K, et al. Work site-based cancer prevention: primary results from the Working Well trial. *Am J Public Health* 1996;86:939-47.



## References

- Soria R, Legido A, Escolano C. A randomised controlled trial of motivational interviewing for smoking cessation. *Br J Gen Pract* 2006;56:768-74.
- Stahre M, Roeber J, Kanny D, et al. Contribution of excessive alcohol consumption to deaths and years of potential life lost in the United States. *Prev Chronic Dis* 2014 Jun 26;11:E109.
- Stevens VJ, Glasgow RE, Toobert DJ, et al. One-year results from a brief, computer assisted intervention to decrease consumption of fat and increase consumption of fruit and vegetables. *Prev Med* 2003;36:594-600.
- Tait RJ, Christensen H. Internet-based interventions for young people with problematic substance use: a systematic review. *Med J Aust* 2010;192:S15-21.
- Task Force on Community Preventive Services. Preventive services: what works to promote health? Oxford University Press. 2005.
- Tessaro I, Rye S, Parker L, et al. Effectiveness of a nutrition intervention with rural low-income women. *Am J Health Behav* 2007;31:35-43.
- Tinker LF, Rosal MC, Young AF, et al. Predictors of dietary change and maintenance in the women's health initiative dietary modification trial. *J Am Diet Assoc* 2007;107:1155-65.
- U.S. Department of Agriculture. Scientific report of the 2015 dietary guidelines advisory committee: advisory report to the secretary of health and human services and the secretary of agriculture. Available at: <http://health.gov/dietaryguidelines/2015-scientific-report/PDFs/Scientific-Report-of-the-2015-Dietary-Guidelines-Advisory-Committee.pdf>. February 2015.
- U.S. Department of Health and Human Services. 2008 physical activity guidelines for Americans. 2008a. (Guideline)
- U.S. Department of Health and Human Services. Dietary guidelines for Americans 2010. Available at: <http://www.dietaryguidelines.gov>. 2010. (Guideline)
- U.S. Department of Health and Human Services. Treating tobacco use and dependence: 2008 update. Available at: <http://www.ahrq.gov/professionals/clinicians-providers/guidelines-recommendations/tobacco/index.html>. 2008b.
- U.S. Department of Health and Human Services – National Institutes of Health. In Your Healthy Guide to Sleep. NIH Publication No. 11-5271. August 2011.
- van Sluijs EMF, van Poppel MNM, van Mechelen W. Stage-based lifestyle interventions in primary care: are they effective? *Am J Prev Med* 2004;26:330-43.
- Vanwormer JJ, Boucher JL, Pronk NP. Telephone-based counseling improves dietary fat, fruit, and vegetable consumption: a best-evidence synthesis. *J Am Diet Assoc* 2006;106:1434-44.
- Vartanian LR, Schwartz MB, Brownell KD. Effects of soft drink consumption on nutrition and health: a systematic review and meta-analysis. *Am J Public Health* 2007;97:667-75.
- Verheijden MW, Van der Veen JE, Bakx JC, et al. Stage-matched nutrition guidance: stages of change and fat consumption in Dutch patients at elevated cardiovascular risk. *J Nutr Educ Behav* 2004;36:228-37.
- Wang CJ, Fetzer SJ, Yang YC, Wang WL. The efficacy of using self-monitoring diaries in a weight loss program for chronically ill obese adults in a rural area. *J Nurs Res* 2012;20:181-88.
- Wang X, Ouyang Y, Liu J, et al. Fruit and vegetable consumption and mortality from all causes, cardiovascular disease, and cancer: systematic review and dose-response meta-analysis of prospective cohort studies. *BMJ* 2014;349:g4490.

**References**

---

[Watson NF, Badr MS, Belenky G, et al.](#) Recommended amount of sleep for a healthy adult: a joint consensus statement of the American academy of sleep medicine and sleep research society. *Sleep* 2015;38:843-44.

[White A, Kavanagh D, Stallman H, et al.](#) Online alcohol interventions: a systematic review. *J Med Internet Res* 2010;12:e62.

Whitlock EP, Orleans CT, Pender N, Allan J. Evaluating primary care behavioral counseling interventions: an evidence-based approach. *J Prev Med* 2002;22:267-84.

[World Health Organization \(WHO\).](#) *In Global Recommendations on Physical Activity for Health.* 2010.