

References

Links are provided for those new references added to this edition (author name is highlighted in blue).

[Abraira C, Duckworth WC, Moritz T](#). Glycaemic separation and risk factor control in the veterans affairs diabetes trial: an interim report. *Diabetes Obes Metab* 2009;11:150-56. (High Quality Evidence)

ACCORD Study Group, The. Effects of combination lipid therapy in type 2 diabetes mellitus. *N Engl J Med* 2010a;362:1563-74. (High Quality Evidence)

ACCORD Study Group, The. Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med* 2010b;362:1575-85. (High Quality Evidence)

[ACCORD Study Group, The](#). Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545-59. (High Quality Evidence)

[Ackermann RT, Cheng YJ, Williamson DF, Gregg EW](#). Identifying adults at high risk for diabetes and cardiovascular disease using hemoglobin A1c national health and nutrition examination survey 2005-2006. *Am J Prev Med* 2011;40:11-17. (Low Quality Evidence)

ADVANCE Collaborative Group, The. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2008;358:2560-72. (High Quality Evidence)

AIM-HIGH Investigators, Boden WE, Probstfield JL, et al. Niacin in patients with low HDL cholesterol levels receiving intensive statin therapy. *N Engl J Med* 2011;365:2255-67. (Moderate Quality Evidence)

[Ajala O, English P, Pinkney J](#). Systematic review and meta-analysis of different dietary approaches to the management of type 2 diabetes. *Am J Clin Nutr* 2013;97:505-16. (Systematic Review)

[Al-Shareef MA, Sanneh AF, Aljoudi AS](#). Clinical effect of metformin in children and adolescents with type 2 diabetes mellitus: a systematic review and meta-analysis. *J Family Community Med* 2012;19:68-73. (Systematic Review)

Alkharouf J, Nalinikumari K, Corry D Tuck M. Long term effects of the angiotensive converting enzyme inhibitor captopril on metabolic control in non-insulin dependent diabetes mellitus. *Am J Hypertension* 1993;6:337-43. (Reference)

ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group, The. Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium channel blocker vs diuretic: the antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT). *JAMA* 2002;288:2981-97. (High Quality Evidence)

American College of Physicians, American Diabetes Association, and American Academy of Ophthalmology. Screening guidelines for diabetic retinopathy. *Ann Intern Med* 1992;116:683-85. (Guideline)

[American Diabetes Association](#). Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2010;33:S62-S69. (Reference)

American Diabetes Association. Hospital admission guidelines for diabetes. *Diabetes Care* 2004a;27:S103. (Reference)

American Diabetes Association. Nephropathy in diabetes. *Diabetes Care* 2004b;27:S79-S83. (Reference)

American Diabetes Association. Peripheral arterial disease in people with diabetes. *Diabetes Care* 2003;26:3333-41. (Guideline)

American Diabetes Association. Physical activity/exercise and diabetes. *Diabetes Care* 2004c;27:S58-S62. (Guideline)

References

- American Diabetes Association. Preventive foot care in diabetes. *Diabetes Care* 2004d;27:S63-S64. (Reference)
- American Diabetes Association. Self monitoring of blood glucose (consensus statement). *Diabetes Care* 1994;17:81-86. (Reference)
- American Diabetes Association. Standards of medical care in diabetes – 2014. *Diabetes Care* 2014;37:S14-80. (Guideline)
- American Diabetes Association. Statistics about diabetes. Available at: <http://www.diabetes.org/diabetes-basics/statistics/>. 2012. (Reference)
- Anderson RJ, Freedland KE, Clouse RE, Lustman PJ. The prevalence of comorbid depression in adults with diabetes: a meta-analysis. *Diabetes Care* 2001;24:1069-78. (Meta-analysis)
- Anderson RT, Narayan KM, Feeney P, et al. Effect of intensive glycemic lowering on health-related quality of life in type 2 diabetes: ACCORD trial. *Diabetes Care* 2011;34:807-12. (High Quality Evidence)
- Andrews RC, Cooper AR, Montgomery AA, et al. Diet or diet plus physical activity versus usual care in patients with newly diagnosed type 2 diabetes: the early ACTID randomised controlled trial. *Lancet* 2011;378:129-39. (High Quality Evidence)
- Antithrombotic Trialists' (ATT) Collaboration. Aspirin in the primary and secondary prevention of vascular disease: collaborative meta-analysis of individual participant data from randomised trials. *Lancet* 2009;373:1849-60. (Meta-analysis)
- Appel LJ, Wright Jr JT, Greene T, et al. Intensive blood-pressure control in hypertensive chronic kidney disease. *N Engl J Med* 2010;363:918-29. (Reference)
- Arguedas JA, Leiva V, Wright JM. Blood pressure targets for hypertension in people with diabetes mellitus. *Cochrane Database Syst Rev* 2013;10:CD008277. (Systematic Review)
- Ash S, Reeves MM, Yeo S, et al. Effect of intensive dietetic interventions on weight and glycaemic control in overweight men with type II diabetes: a randomised trial. *Int J Obes Relat Metab Disord* 2003;27:797-802. (Moderate Quality Evidence)
- Azadbakht L, Fard NR, Karimi M, et al. Effects of the dietary approaches to stop hypertension (DASH) eating plan on cardiovascular risks among type 2 diabetic patients: a randomized crossover clinical trial. *Diabetes Care* 2011;34:55-57. (High Quality Evidence)
- Bangalore S, Kumar S, Lobach I, Messerli FH. Blood pressure targets in subjects with type 2 diabetes mellitus/impaired fasting glucose: observations from traditional and bayesian random-effects meta-analyses of randomized trials. *Circulation* 2011;123:2799-2810. (Meta-analysis)
- Belch J, MacCuish A, Campbell I, et al. The prevention of progression of arterial disease and diabetes (POPADAD) trial: factorial randomised placebo controlled trial of aspirin and antioxidants in patients with diabetes and asymptomatic peripheral arterial disease. *BMJ* 2008;337:a1840. (High Quality Evidence)
- Bennett PH, Haffner S, Kasiske BL, et al. Diabetic renal disease recommendations: screening and management of microalbuminuria in patients with diabetes mellitus: recommendations to the Scientific Advisory Board of the National Kidney Foundation from an ad hoc committee of the Council on Diabetes Mellitus of the National Kidney Foundation. *Am J Kidney Dis* 1995;25:107-35. (Reference)
- Boulé NG, Haddad E, Kenny GP, et al. Effects of exercise on glycemic control and body mass in type 2 diabetes mellitus: a meta-analysis of controlled clinical trials. *JAMA* 2001;286:1218-27. (Meta-analysis)
- Brehm BJ, Lattin BL, Summer SS, et al. One-year comparison of a high-monosaturated fat diet with a high-carbohydrate diet in type 2 diabetes. *Diabetes Care* 2009;32:215-20. (High Quality Evidence)

References

- Brunerova L, Smejkalova V, Potockova J, Andel M. A comparison of the influence of a high-fat diet enriched in monounsaturated fatty acids and conventional diet on weight loss and metabolic parameters in obese non-diabetic and type 2 diabetic patients. *Diabet Med* 2007;24:533-40. (Moderate Quality Evidence)
- Buchwald H, Estok R, Fahrbach K, et al. Weight and type 2 diabetes after bariatric surgery: systematic review and meta-analysis. *Am J Med* 2009;122:248-56. (Systematic Review)
- Callaghan BC, Little AA, Feldman EL, Hughes RAC. Enhanced glucose control for preventing and treating diabetic neuropathy. *Cochrane Database Syst Rev* 2012;6:CD007543. (Systematic Review)
- Campbell CL, Smyth S, Montalescot G, Steinhubl SR. Aspirin dose for the prevention of cardiovascular disease: a systematic review. *JAMA* 2007;297:2018-24. (Systematic Review)
- Cannon CP, Braunwald E, McCabe CH, et al. Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med* 2004;350:1495-504. (High Quality Evidence)
- Casagrande SS, Cowie CC, Fradkin JE. Utility of the U.S. preventive services task force criteria for diabetes screening. *Am J Prev Med* 2013;45:167-74. (Low Quality Evidence)
- Castaneda C, Layne JE, Munoz-Orians L, et al. A randomized controlled trial of resistance exercise training to improve glycemic control in older adults with type 2 diabetes. *Diabetes Care* 2002;25:2335-41. (High Quality Evidence)
- Cauza E, Hanusch-Enserer U, Strasser B, et al. The relative benefits of endurance and strength training on the metabolic factors and muscle function of people with type 2 diabetes mellitus. *Arch Phys Med Rehabil* 2005;86:1527-33. (Low Quality Evidence)
- Centers for Disease Control, The. National diabetes fact sheet, 2011: fast facts on diabetes. 2011. (Reference)
- Chiasson JL, Josse RG, Gomis R, et al. Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomised trial. *Lancet* 2002;359:2072-77. (Reference)
- Cholesterol Treatment Trialists' (CTT) Collaboration. Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials. *Lancet* 2010;376:1670-81. (Meta-analysis)
- Cholesterol Treatment Trialists' (CTT) Collaborators. Efficacy of cholesterol-lowering therapy in 18,686 people with diabetes in 14 randomised trials of statins: a meta-analysis. *Lancet* 2008;371:117-25. (Meta-analysis)
- Church TS, Blair SN, Cocreham S, et al. Effects of aerobic and resistance training on hemoglobin A1c levels in patients with type 2 diabetes: a randomized controlled trial. *JAMA* 2010;304:2253-62. (High Quality Evidence)
- Clement S, Braithwaite SS, Magee MF, et al. Management of diabetes and hyperglycemia in hospitals. *Diabetes Care* 2004;27:553-91. (Reference)
- Cohen RV, Pinheiro JC, Schiavon CA, et al. Effects of gastric bypass surgery in patients with type 2 diabetes and only mild obesity. *Diabetes Care* 2012;35:1420-28. (Low Quality Evidence)
- Colosia AD, Palencia R, Khan S. Prevalence of hypertension and obesity in patients with type 2 diabetes mellitus in observational studies: a systematic literature review. *Diabetes Metab Syndr Obes* 2013;6:327-38. (Low Quality Evidence)
- Cowie CC, Rust KF, Byrd-Holt DD, et al. Prevalence of diabetes and high risk for diabetes using A1c criteria in the U.S. population in 1988-2006. *Diabetes Care* 2010;33:562-68. (Low Quality Evidence)

References

- Cryer PE, Davis SN, Shamoon H. Hypoglycemia in diabetes. *Diabetes Care* 2003;26:1902-12. (Reference)
- Davidson PC, Hebblewhite HR, Bode BW, et al. Statistically based CSII parameters: correction factor (CF) (1700 rule), carbohydrate-insulin ratio (CIR) (2.8 rule), and basal-to-total ratio. *Diabetes Technol Ther* 2003;5:237. (Reference)
- Davidson JK. Endogenous insulin, proinsulin, and C-peptide. In Clinical Diabetes Mellitus: A Problem Oriented Approach. Thieme Inc., New York. 1986;219-22. (Reference)
- DeFronzo RA. Diabetic nephropathy: etiologic and therapeutic considerations. *Diabetes Reviews* 1995;3:510-64. (Reference)
- Deakin TA, McShane CE, Cade JE, Williams RD. Group based training for self-management strategies in people with type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2009;(2):CD003417. (Systematic Review)
- de Groot M, Anderson R, Freedland KE, et al. Association of depression and diabetes complications: a meta-analysis. *Psychosom Med* 2001;63:619-30. (Reference)
- Diabetes Prevention Program Research Group. 10-year follow-up of diabetes incidence and weight loss in the diabetes prevention program outcomes study. *Lancet* 2009;374:1677-86. (High Quality Evidence)
- Diabetic Retinopathy Study Research Group, The. Photocoagulation treatment of proliferative diabetic retinopathy: clinical application of diabetic retinopathy, (DRS) findings, DRS report number 8. *Ophthalmology* 1981;88:583-600. (Reference)
- Dorman RB, Serrot FJ, Miller CJ, et al. Case-matched outcomes in bariatric surgery for treatment of type 2 diabetes in the morbidly obese patient. *Ann Surg* 2012;255:287-93. (Low Quality Evidence)
- Droumaguet C, Balkau B, Simon D, et al. Use of HbA1c in predicting progression to diabetes in French men and women: data from an epidemiological study on the insulin resistance syndrome (DESIR). *Diabetes Care* 2006;29:1619-25. (Low Quality Evidence)
- Duckworth W, Abraira C, Moritz T, et al. Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med* 2009;360:129-39. (High Quality Evidence)
- Duke SAS, Colagiuri S, Colagiuri R. Individual patient education for people with type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2009;(1):CD005268. (Systematic Review)
- Duncan I, Ahmed T, Li Q, et al. Assessing the value of the diabetes educator. *Diabetes Educ* 2011;37:638-57. (Cost-Effectiveness)
- Dunstan DW, Daly RM, Owen N, et al. High-intensity resistance training improves glycemic control in older patients with type 2 diabetes. *Diabetes Care* 2002;25:1729-36. (High Quality Evidence)
- Elhayany A, Lustman A, Abel R, et al. A low carbohydrate Mediterranean diet improves cardiovascular risk factors and diabetes control among overweight patients with type 2 diabetes mellitus: a 1-year prospective randomized intervention study. *Diabetes Obes Metab* 2010;12:204-09. (High Quality Evidence)
- Eriksson J, Lindstrom J, Valle T, et al. Prevention of type II diabetes in subjects with impaired glucose tolerance: the diabetes prevention study (DPS) in Finland. *Diabetologia* 1999;42:793-801. (Reference)
- Esposito K, Maiorino MI, Ciotola M, et al. Effects of a Mediterranean-style diet on the need for antihyperglycemic drug therapy in patients with newly diagnosed type 2 diabetes: a randomized trial. *Ann Intern Med* 2009;151:306-14. (Moderate Quality Evidence)

References

- Estacio RO, Coll JR, Tran ZV, Schrier RW. Effect of intensive blood pressure control with valsartan on urinary albumin excretion in normotensive patients with type 2 diabetes. *Am J Hypertens* 2006;19:1241-48. (High Quality Evidence)
- Estruch R, Ros E, Salas-Salvadó J, et al. Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med* 2013;368:1279-90. (High Quality Evidence)
- ETDRS Research Group. Early photocoagulation for diabetic retinopathy: ETDRS report number 9. *Ophthalmology* 1991;98:766-85. (Reference)
- ETDRS Research Group. Photocoagulation for diabetic macular edema: ETDRS report number 1. *Arch Ophthalmol* 1985;103:1796-1806. (Reference)
- Evert AB, Boucher JL, Cypress M, et al. Nutrition therapy recommendations for the management of adults with diabetes. *Diabetes Care* 2014;37:S120-43. (Guideline)
- Fong DS, Aiello L, Gardner TW, et al. Retinopathy in diabetes. *Diabetes Care* 2004;27:S84-S87. (Reference)
- Foster GD, Borradaile KE, Sanders MH, et al. A randomized study on the effect of weight loss on obstructive sleep apnea among obese patients with type 2 diabetes: the sleep AHEAD study. *Arch Intern Med* 2009a;169:1619-26. (Reference)
- Foster GD, Sanders MH, Millman R, et al. Obstructive sleep apnea among obese patients with type 2 diabetes. *Diabetes Care* 2009b;32:1017-19. (Reference)
- Furnary AP, Braithwaite SS. Effects of outcome on in-hospital transition from intravenous insulin infusion to subcutaneous therapy. *Am J Cardiol* 2006;98:557-64. (Reference)
- Gaede P, Lund-Andersen H, Parving HH, Pedersen O. Effect of a multifactorial intervention on mortality in type 2 diabetes. *N Engl J Med* 2008;358:580-91. (High Quality Evidence)
- Gaede P, Vedel P, Larsen N, et al. Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *N Engl J Med* 2003;348:383-93. (High Quality Evidence)
- Garber AJ, Moghissi ES, Bransome Jr ED, et al. American college of endocrinology position statement on inpatient diabetes and metabolic control. *Endocr Pract* 2004;10:77-82. (Reference)
- Gary TL, Genkinger JM, Guallar E, et al. Meta-analysis of randomized educational and behavioral interventions in type 2 diabetes. *Diabetes Educ* 2003;29:488-501. (Meta-analysis)
- Gerich JE. Novel insulins: expanding options in diabetes management. *Am J Med* 2002;113:339-40. (Reference)
- Gibbs BB, Brancati FL, Chen H, et al. Effect of improved fitness beyond weight loss on cardiovascular risk factors in individuals with type 2 diabetes in the Look AHEAD study. *Eur J Prev Cardiol* 2014;21:608-17. (High Quality Evidence)
- Gillett M, Dallosso HM, Dixon S, et al. Delivering the diabetes education and self management for ongoing and newly diagnosed (DESMOND) programme for people with newly diagnosed type 2 diabetes: cost effectiveness analysis. *BMJ* 2010;341:c4093. (Cost-Effectiveness Analysis)
- Gregg EW, Cadwell BL, Cheng YJ, et al. Trends in the prevalence and ratio of diagnosed to undiagnosed diabetes according to obesity levels in the U.S. *Diabetes Care* 2004;27:2806-12. (Low Quality Evidence)
- Guerci B, Sauvanet JP. Subcutaneous insulin: pharmacokinetic variability and glycemic variability. *Diabetes Metab* 2005;31:4S7-4S24. (Reference)

References

- Guicciardi M, Lecis R, Anziani C, et al. Type 2 diabetes mellitus, physical activity, exercise self-efficacy, and body satisfaction: an application of the transtheoretical model in older adults. *Health Psychol and Behav Med* 2014;2:748-58. (Low Quality Evidence)
- Hannah R, Levin N, London R, et al, eds. Renal disease in the managed care setting: selection and monitoring of outcome criteria. *Am J Kidney Dis* 1999;33(Suppl 1):S1-S23. (Reference)
- Hansson L, Zanchetti A, Carruthers SG, et al. Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the hypertension optimal treatment (HOT) randomised trial. *Lancet* 1998;361:1755-62. (Moderate Quality Evidence)
- He FJ, Li J, MacGregor GA. Effect of longer-term modest salt reduction on blood pressure. *Cochrane Database Syst Rev* 2013;4:CD004937. (Systematic Review)
- Heart Outcomes Prevention Evaluation Study Investigators, The. Effects of an angiotensin-converting – enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. *N Engl J Med* 2002;342:145-53. (Reference)
- Heart Outcomes Prevention Evaluation (HOPE) Study Investigators. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. *Lancet* 2000;355:253-59. (High Quality Evidence)
- Heart Protection Study Collaborative Group. MRC/BHF heart protection study of cholesterol lowering with simvastatin in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet* 2002; 360:7-22. (High Quality Evidence)
- Heart Protection Study Collaborative Group. MRC/BHF heart protection study of cholesterol-lowering with simvastatin in 5,963 people with diabetes: a randomised placebo-controlled trial. *Lancet* 2003;361:2005-16. (High Quality Evidence)
- Hemmingsen B, Lund SS, Gluud C, et al. Targeting intensive glycaemic control versus targeting conventional glycaemic for type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2013;11:CD008143. (Systematic Review)
- Hirsch IB. Insulin in the hospital setting. New York, Adelphi, 2002;25. (Reference)
- Hirsch IB, Paauw DS, Brunzell J, et al. Inpatient management of adults with diabetes. *Diabetes Care* 1995;18:870. (Reference)
- Hoerger TJ, Zhang P, Segel JE, et al. Cost-effectiveness of bariatric surgery for severely obese adults with diabetes. *Diabetes Care* 2010;33:1933-39. (Low Quality Evidence)
- Holman RR, Farmer AJ, Davies MJ, et al. Three-year efficacy of complex insulin regimens in type 2 diabetes. *N Engl J Med* 2009;361:1736-47. (Reference)
- Holman RR, Paul SK, Bethel MA, et al. Long-term follow-up after tight control of blood pressure in type 2 diabetes. *N Engl J Med* 2008a;359:1565-76. (High Quality Evidence)
- Holman RR, Paul SK, Bethel MA, et al. 10-year follow-up of intensive glucose control in type 2 diabetes. *N Engl J Med* 2008b;359:1577-89. (Low Quality Evidence)
- Howard BV, Roman MJ, Devereux RB, et al. Effect of lower targets for blood pressure and LDL cholesterol on atherosclerosis in diabetes: the SANDS randomized trial. *JAMA* 2008;299:1678-89. (High Quality Evidence)
- Ikramuddin S, Korner J, Lee WJ, et al. Roux-en-Y gastric bypass vs intensive medical management for the control of type 2 diabetes, hypertension, and hyperlipidemia: the diabetes surgery study randomized clinical trial. *JAMA* 2013;309:2240-49. (High Quality Evidence)

References

- Institute of Medicine. Sodium intake in populations: assessment of evidence. May 2013. (Reference)
- International Expert Committee, The. International expert committee report on the role of A1c assay in the diagnosis of diabetes. *Diabetes Care* 2009;32:1327-34. (Low Quality Evidence)
- Ismail-Beigi F, Craven T, Banerji MA, et al. Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. *Lancet* 2010;376:419-30. (High Quality Evidence)
- James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the eighth joint national committee (JNC 8). *JAMA* 2014;311:507-20. (Reference)
- John R, Fogelfeld L. Inpatient management of diabetes and hyperglycemia. *Dis Mon* 2004;50:438-79. (Reference)
- Karter AJ, Ferrara A, Liu JY, et al. Ethnic disparities in diabetic complications in an insured population. *JAMA* 2002;287:2519-27. (Low Quality Evidence)
- Kelley DE, Bray GA, Pi-Sunyer FX, et al. Clinical efficacy of orlistat therapy in overweight and obese patients with insulin-treated type 2 diabetes. *Diabetes Care* 2002;25:1033-41. (Reference)
- Klein R, Klein BEK, Moss SE, et al. The Wisconsin epidemiologic study of diabetic retinopathy. III. Prevalence and risk of diabetic retinopathy when age at diagnosis is 30 or more years. *Arch Ophthalmol* 1984;102:527-32. (Reference)
- Klein R, Moss SE, Klein BEK. New management concepts for timely diagnosis of diabetic retinopathy treatable by photocoagulation. *Diabetes Care* 1987;10:633-38. (Reference)
- Kumar PR, Bhansali A, Ravikiran M, et al. Utility of glycated hemoglobin in diagnosing type 2 diabetes mellitus: a community-based study. *J Clin Endocrinol Metab* 2010;95:2832-35. (Moderate Quality Evidence)
- Lakerveld J, Bot SD, Chinapaw MJ, et al. Motivational interviewing and problem solving treatment to reduce type 2 diabetes and cardiovascular disease risk in real life: a randomized controlled trial. *Int J Behav Nutr Phys Act* 2013;10:47. (Moderate Quality Evidence)
- Lewis EJ, Hunsicker LG, Bain RP, et al. The effect of angiotensin-converting-enzyme inhibition on diabetic nephropathy. *N Engl J Med* 1993;329:1456-62. (Reference)
- Lewis EJ, Hunsicker LG, Clarke WR, et al. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001;345:851-60. (Reference)
- Lewis JB. Blood pressure control in chronic kidney disease: is less really more? *J Am Soc Nephrol* 2010;21:1086-92. (Reference)
- Li C, Lai Y, Tseng C, et al. A population study on the association between leisure time physical activity and self-rated health among diabetics in Taiwan. *BMC Public Health* 2010;10:277. (Moderate Quality Evidence)
- Li G, Zhang P, Wang J, et al. The long-term effect of lifestyle interventions to prevent diabetes in the China Da Qing diabetes prevention study: a 20-year follow-up study. *Lancet* 2008;371:1783-89. (Low Quality Evidence)
- Lipska KJ, Bailey CJ, Inzucchi SE. Use of metformin in the setting of mild-to-moderate renal insufficiency. *Diabetes Care* 2011;34:1431-37. (Low Quality Evidence)

References

- Look AHEAD Research Group, The. Cardiovascular effects of intensive lifestyle intervention in type 2 diabetes. *N Engl J Med* 2013;369:145-54. (High Quality Evidence)
- Lustman PJ, Gavard JA. Psychosocial aspects of diabetes in adult populations. *In Diabetes in America*, 2nd Ed. 507-18. On-line reference, accessed July, 2001 at <http://diabetes-in-america.s-3.com/>. (Reference)
- Macchia A, Laffaye N, Comignani PD, et al. Statins but not aspirin reduce thrombotic risk assessed by thrombin generation in diabetic patients without cardiovascular events: the RATIONAL trial. *PLoS* 2012;7:e32894. (High Quality Evidence)
- Maciejewski ML, Livingston EH, Smith VA, et al. Survival among high-risk patients after bariatric surgery. *JAMA* 2011;305:2419-26. (Low Quality Evidence)
- Makary MA, Clarke JM, Shore AD, et al. Medication utilization and annual health care costs in patients with type 2 diabetes mellitus before and after bariatric surgery. *Arch Surg* 2010;145:726-31. (Low Quality Evidence)
- Mayfield JA, Reiber GE, Sanders LJ, et al. Preventive foot care in people with diabetes. *Diabetes Care* 1998;21:2161-77. (Reference)
- Miller CK, Edwards L, Kissling G, Sanville L. Nutrition education improves metabolic outcomes among older adults with diabetes mellitus: results from a randomized controlled trial. *Prev Med* 2002;34:252-59. (Moderate Quality Evidence)
- Mingrone G, Panunzi S, De Gaetano A, et al. Bariatric surgery versus conventional medical therapy for type 2 diabetes. *N Engl J Med* 2012;366:1577-85. (Moderate Quality Evidence)
- Mogensen CE, Keane WF, Bennett PH, et al. Prevention of diabetic renal disease with special reference to microalbuminuria. *Lancet* 1996;346:1080-84. (Reference)
- Morrish NJ, Stevens LK, Fuller JH, et al. Risk factors for macrovascular disease in diabetes mellitus: the London follow-up to the WHO multinational study of vascular disease in diabetics. *Diabetologia* 1991;34:590-94. (Low Quality Evidence)
- National Institutes of Health. Morbidity and mortality of dialysis (consensus statement, online). 1993;11:1-33. (Reference)
- Nelson RG, Knowler WC, Pettitt DJ, et al. Assessment of risk of overt nephropathy in diabetic patients from albumin excretion in untimed urine specimens. *Arch Intern Med* 1991;151:1761-65. (Reference)
- NICE-SUGAR Study Investigators, The. Intensive versus conventional glucose control in critically ill patients. *N Engl J Med* 2009;360:1283-97. (High Quality Evidence)
- Nichols-English G, Poirier S. Optimizing adherence to pharmaceutical care plans. *J Am Pharm Assoc* 2000;40:475-83. (Reference)
- Nield L, Moore HJ, Hooper L, et al. Dietary advice for treatment of type 2 diabetes mellitus in adults. *Cochrane Database Syst Rev* 2007;(3):CD004097. (Systematic Review)
- Nilsson PM. Target blood pressure in diabetes patients with hypertension – what is the accumulated evidence in 2011? *J Zhejiang Univ Sci B* 2011;12:611-23. (Systematic Review)
- Norris SL, Zhang X, Avenell A, et al. Long-term non-pharmacological weight loss interventions for adults with type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2005;(2):CD004095. (Systematic Review)
- O'Connor PJ. Commentary – improving diabetes care by combating clinical inertia. *Health Serv Res* 2005a;40:1854-61. (Low Quality Evidence)
- O'Connor PJ. Overcome clinical inertia to control systolic blood pressure. *Arch Intern Med* 2003;163:2677-78. (Low Quality Evidence)

References

- O'Connor PJ, Crabtree BF, Yanoshik MK. Differences between diabetic patients who do and do not respond to a diabetes care intervention: a qualitative analysis. *Fam Med* 1997;29:424-28. (Low Quality Evidence)
- O'Connor PJ, Sperl-Hillen JM, Johnson PE, et al. Advances in patient safety: clinical inertia and outpatient medical errors. *AHRQ* 2005b;2:293-308. (Low Quality Evidence)
- Ogawa H, Nakayama M, Morimoto T, et al. Low-dose aspirin for primary prevention of atherosclerotic events in patients with type 2 diabetes: a randomized controlled trial. *JAMA* 2008;300:2134-41. (High Quality Evidence)
- Olson DE, Rhee MK, Herrick K, et al. Screening for diabetes and pre-diabetes with proposed A1c-based diagnostic criteria. *Diabetes Care* 2010;33:2184-89. (Low Quality Evidence)
- Orozco LJ, Buchleitner AM, Gimenez-Perez G, et al. Exercise or exercise and diet for preventing type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2008;(3):CD003054. (Systematic Review)
- Pal K, Eastwood SV, Michie S, et al. Computer-based diabetes self-management interventions for adults with type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2013;3:CD008776. (Systematic Review)
- Pettitt DJ, Opsina P, Kolaczynski JW, Jovanovic L. Comparison of an insulin analog, insulin aspart, and regular human insulin with no insulin in gestational diabetes mellitus. *Diabetes Care* 2003;26:183-86. (Reference)
- Pignone M, Eamshaw S, Tice J, Pletcher MJ. Aspirin, statins, or both drugs for the primary prevention of coronary heart disease events in men: a cost-utility analysis. *Ann Intern Med* 2006;144:326-36 (Low Quality Evidence)
- Polonsky WH, Fisher L, Schikman CH, et al. A structured self-monitoring of blood glucose approach in type 2 diabetes encourages more frequent, intensive, and effective physician interventions: results from the STeP study. *Diabetes Technol Ther* 2011;13:797-802. (Reference)
- Polonsky KS, Given BD, Hirsch ET, et al. Quantitative study of insulin secretion and clearance in normal and obese subjects. *J Clin Invest* 1988a;81:435-41. (Reference)
- Polonsky KS, Given BD, Van Cauter E. Twenty-four-hour profiles and pulsatile patterns of insulin secretion in normal and obese subjects. *J Clin Invest* 1988b;81:442-48. (Reference)
- Radhakrishnan K. The efficacy of tailored interventions for self-management outcomes of type 2 diabetes, hypertension or heart disease: a systematic review. *J Adv Nurs* 2011;68:496-510. (Systematic Review)
- Rahman M, Simmons RK, Hennings SH, et al. Effect of screening for type 2 diabetes on population-level self-rated health outcomes and measures of cardiovascular risk: 13-year follow-up of the Ely cohort. *Diabet Med* 2012;29:886-92. (Low Quality Evidence)
- Rave K, Klein O, Frick AD, Becker RHA. Advantage of premeal-injected insulin glulisine compared with regular human insulin in subjects with type 1 diabetes. *Diabetes Care* 2006;29:1812-17. (Reference)
- Ravid M, Savin H, Jutrin I, et al. Long-term stabilizing effect of angiotensin-converting enzyme inhibition on plasma creatinine and on proteinuria in normotensive type II diabetic patients. *Ann Int Med* 1993;118:577-81. (Moderate Quality Evidence)
- Ray KK, Seshasai SR, Wijesuriya S, et al. Effect of intensive control of glucose on cardiovascular outcomes and death in patients with diabetes mellitus: a meta-analysis of randomised controlled trials. *Lancet* 2009;373:1765-72. (Meta-analysis)
- Reynolds NA, Wagstaff AJ. Insulin aspart: a review of its use in the management of type 1 or type 2 diabetes mellitus. *Drugs* 2004;64:1957-74. (Reference)

References

- Robbins JM, Thatcher GE, Webb DA, Valdmanis VG. Nutritionist visits, diabetes classes, and hospitalization rates and charges: the urban diabetes study. *Diabetes Care* 2008;31:655-60. (Low Quality Evidence)
- Rosiak M, Postula M, Kaplon-Cieslicka A, et al. The effect of doubling the dose of acetylsalicylic acid (ASA) on platelet function parameters in patients with type 2 diabetes and platelet hyperreactivity during treatment with 75 mg of ASA: a subanalysis of the AVOCADO study. *Kardiol Pol* 2013;71:552-57. (Moderate Quality Evidence) (Moderate Quality Evidence)
- Saenz A, Fernandez-Esteban I, Mataix A, et al. Metformin monotherapy for type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2005;(3):CD002966. (Systematic Review)
- Salpeter SR, Greyber E, Pasternak GA, Salpeter EE. Risk of fatal and nonfatal lactic acidosis with metformin use in type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2010;(4):CD002967. (Systematic Review)
- Schauer PR, Kashyap SR, Wolski K, et al. Bariatric surgery versus intensive medical therapy in obese patients with diabetes. *N Engl J Med* 2012;366:1567-76. (High Quality Evidence)
- Selvin E, Bolen S, Yeh HC, et al. Cardiovascular outcomes in trials of oral diabetes medications: a systematic review. *Arch Intern Med* 2008;168:2070-80. (High Quality Evidence)
- Sigal RJ, Kenny GP, Wasserman DH, Castaneda-Sceppa C. Physical activity/exercise and type 2 diabetes. *Diabetes Care* 2004;27:2518-39. (Low Quality Evidence)
- Siminerio LM. Implementing diabetes self-management training programs: breaking through the barriers in primary care. *Endocr Pract* 2006a;12:124-30. (Low Quality Evidence)
- Siminerio LM, Piatt G, Zgibor JC. Implementing the chronic care model for improvements in diabetes care and education in a rural primary care practice. *Diabetes Educ* 2005;31:225-34. (Low Quality Evidence)
- Siminerio LM, Piatt GA, Emerson S, et al. Deploying the chronic care model to implement and sustain diabetes self-management training programs. *Diabetes Educ* 2006b;32:253-60. (Low Quality Evidence)
- Simmons RK, Echouffo-Tcheugui JB, Sharp SJ, et al. Screening for type 2 diabetes and population mortality over 10 years (ADDITION-Cambridge): a cluster-randomized controlled trial. *Lancet* 2012;380:1741-48. (High Quality Evidence)
- Soejima H, Ogawa H, Morimoto T, et al. Aspirin reduces cerebrovascular events in type 2 diabetic patients with poorly controlled blood pressure. *Circ J* 2012;76:1526-32. (Low Quality Evidence)
- Sperl-Hillen JM, O'Connor JP. Factors driving diabetes care improvement in a large medical group: ten years of progress. *Am J Manag Care* 2005;11:S177-S185. (Low Quality Evidence)
- Steinsbekk A, Rygg LO, Lisulo M, et al. Group based diabetes self-management education compared to routine treatment for people with type 2 diabetes mellitus: a systematic review with meta-analysis. *BMC Health Serv Res* 2012;12:213. (Systematic Review)
- Stone NJ, Robinson JG, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American college of cardiology/American heart association task force on practice guidelines. *Circulation* 2014;129:S1-S45. (Reference)
- Suckling RJ, He FJ, MacGregor GA. Altered dietary salt intake for preventing and treating diabetic kidney disease. *Cochrane Database Syst Rev* 2010(12):CD006763. (Systematic Review)
- Taylor F, Huffman MD, Macedo AF, et al. Statins for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev* 2013;1:CD004816. (Systematic Review)
- Thomas D, Elliott EJ. Low glycaemic index, or low glycaemic load, diets for diabetes mellitus. *Cochrane Database Syst Rev* 2009;(1):CD006296. (Systematic Review)

References

- Thomas DE, Elliott EJ, Naughton GA. Exercise for type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2006;(3):CD002968. (Systematic Review)
- Thorpe CT, Fahey LE, Johnson H, et al. Facilitating healthy coping in patients with diabetes: a systematic review. *Diabetes Educ* 2013;39:33-52. (Systematic Review)
- Tricco AC, Ivers NM, Grimshaw JM, et al. Effectiveness of quality improvement strategies on the management of diabetes: a systematic review and meta-analysis. *Lancet* 2012;379:2252-61. (Systematic Review)
- Tshiananga JK, Kocher S, Weber C, et al. The effect of nurse-led diabetes self-management education on glycosylated hemoglobin and cardiovascular risk factors: a meta-analysis. *Diabetes Educ* 2012;38:108-23. (Meta-analysis)
- Tuomilehto J, Lindström J, Eriksson JG, et al. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 2001;344:1343-50. (High Quality Evidence)
- Turnbull FM, Abraira C, Anderson RJ, et al. Intensive glucose control and macrovascular outcomes in type 2 diabetes. *Diabetologia* 2009;52:2288-98. (Meta-analysis)
- UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998a;352:837-53. (High Quality Evidence)
- UK Prospective Diabetes Study (UKPDS) Group. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. *BMJ* 1998b;317:703-13. (High Quality Evidence)
- Umpierrez GE, Isaacs SD, Bazargan N, et al. Hyperglycemia: an independent marker of in-hospital mortality in patients with undiagnosed diabetes. *J Clin Endocrinol Metab* 2002;87:978-82. (Reference)
- Umpierrez GE, Smiley D, Zisman A, et al. Randomized study of basal-bolus insulin therapy in the inpatient management of patients with type 2 diabetes (RABBIT 2 trial). *Diabetes Care* 2007;30:2181-86. (Reference)
- U.S. Preventive Services Task Force. Screening for type 2 diabetes mellitus. Available at: <http://www.uspreventiveservicestaskforce.org/Page/Topic/recommendation-summary/diabetes-mellitus-type-2-in-adults-screening?ds=1&s=>. 2008. (Systematic Review)
- Vague P, Selam JL, Skeie S, et al. Insulin Detemir is associated with more predictable glycemic control and reduced risk of hypoglycemia than NPH insulin in patients with type 1 diabetes on a basal-bolus regimen with premeal insulin aspart. *Diabetes Care* 2003;26:590-96. (Reference)
- Valentine N, Dan de Laar FA, van Driel ML. Adenosine-diphosphate (ADP) receptor antagonists for the prevention of cardiovascular disease in type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2012;11:CD005449. (Systematic Review)
- Van den Berghe G, Wouters P, Weekers F, et al. Intensive insulin therapy in critically ill patients. *N Engl J Med* 2001;345:1359-67. (Reference)
- Viberti G. Outcome variables in the assessment of progression of diabetic kidney disease. *Kidney Intl* 1994;45:S121-24. (Reference)
- Wang F, Carabino JM, Vergara CM. Insulin glargine: a systematic review of a long-acting insulin analogue. *Clin Ther* 2003;25:1541-77. (Reference)
- Waugh NR, Shyangdan D, Taylor-Phillips S, et al. Screening for type 2 diabetes: a short report for the national screening committee. *Health Technol Assess* 2013;17:1-90. (Systematic Review)

References

Weinger K, Beverly EA, Lee Y, et al. The effect of a structured behavioral intervention on poorly controlled diabetes: a randomized controlled trial. *Arch Intern Med* 2011;171:1990-99. (Moderate Quality Evidence)

[Wheeler ML](#), [Dunbar SA](#), [Jaacks LM](#), et al. Macronutrients, food groups, and eating patterns in the management of diabetes: a systematic review of the literature, 2010. *Diabetes Care* 2012;35:434-45. (Systematic Review)

[Wiebe N](#), [Padwal R](#), [Field C](#), et al. A systematic review on the effect of sweeteners on glycemic response and clinically relevant outcomes. *BMC Medicine* 2011;9:123. (Systematic Review)

Wing LMH, Reid CM, Ryan P, et al. A comparison of outcomes with angiotensin-converting-enzyme inhibitors and diuretics for hypertension in the elderly. *N Engl J Med* 2003;348:583-92. (Moderate Quality Evidence)

Yki-Järvinen H, Dressler A, Ziemer M, The HOE 901/3002 Study Group. Less nocturnal hypoglycemia and better post-dinner glucose control with bedtime insulin Glargine compared with bedtime NPH insulin during insulin combination therapy in type 2 diabetes. *Diabetes Care* 2000;23:1130-36. (Reference)

[Zoungas S](#), [de Galan BE](#), [Ninomiya T](#), et al. Combined effects of routine blood pressure lowering and intensive glucose control on macrovascular and microvascular outcomes in patients with type 2 diabetes: new results from the ADVANCE trial. *Diabetes Care* 2009;32:2068-74. (High Quality Evidence)