

The subdivisions of this section are:

- References
- Appendices

References

Links are provided for those new references added to this edition (author name is highlighted in [blue](#)).

Accurso AJ, Rastegar DA. The effect of a payer-mandated decrease in buprenorphine dose on aberrant drug tests and treatment retention among patients with opioid dependence. *J Subst Abuse Treat* 2016;61:74-79.

Achterberg WP, Scherder E, Pot AM, Ribbe MW. Cardiovascular risk factors in cognitively impaired nursing home patients: a relationship with pain? *Eur J Pain* 2007;11:707-10.

Adachi T, Fujino H, Nakae A, et al. A meta-analysis of hypnosis for chronic pain problems: a comparison between hypnosis, standard care, and other psychological interventions. *Int J Clin Exp Hypn* 2014;62:1-28.

Afari N, Ahumada SM, Wright LJ, et al. Psychological trauma and functional somatic syndromes: a systematic review and meta-analysis. *Psychosom Med* 2014;76:2-11.

Alam A, Gomes T, Zheng H, et al. Long-term analgesic use after low-risk surgery: a retrospective cohort study. *Arch Intern Med* 2012;172:425-30.

Albert S, Brason FW 2nd, Sanford CK, et al. Project Lazarus: community-based overdose prevention in rural North Carolina. *Pain Med* 2011;12:S77-85.

Alford DP, Compton P, Samet JH. Acute pain management for patients receiving maintenance methadone or buprenorphine therapy. *Ann Intern Med* 2006;144:127-34.

American Geriatrics Society 2015 Beers Criteria Update Expert Panel. American geriatrics society 2015 updated beers criteria for potentially inappropriate medication use in older adults. *J Am Geriatr Soc* 2015;63:2227-46.

American Geriatrics Society Panel on Exercise and Osteoarthritis. Exercise prescription for older adults with osteoarthritis pain: consensus practice recommendations. *JAGS* 2001;49:808-23.

American Pain Society. *In Principles of Analgesic Use in the Treatment of Acute Pain and Cancer Pain*. Sixth Edition. 2008.

American Psychiatric Association. *In Diagnostic and Statistical Manual of Mental Disorders (DSM-5), Fifth Edition*. 2013.

Argoff CE, Kahan M, Sellers EM. Preventing and managing aberrant drug-related behavior in primary care: systematic review of outcomes evidence. *J Opioid Manag* 2014;10:119-34.

Arnold LM, Lu Y, Crofford LJ, et al. A double-blind, multicenter trial comparing duloxetine with placebo in the treatment of fibromyalgia patients with or without major depressive disorder. *Arthritis Rheum* 2004;50:2974-84.

Arnold RM, Han PK, Seltzer D. Opioid contracts in chronic nonmalignant pain management: objectives and uncertainties. *Am J Med* 2006;119:292-96.

Arnow BA. Relationships between childhood maltreatment, adult health and psychiatric outcomes, and medical utilization. *J Clin Psychiatry* 2004;65:10-15.

Asmundson GJ, Coons MJ, Taylor S, Katz J. PTSD and the experience of pain: research and clinical implications of shared vulnerability and mutual maintenance models. *Can J Psychiatry* 2002;47:930-37.

Asmundson GJ, Katz J. Understanding the co-occurrence of anxiety disorders and chronic pain: state-of-the-art. *Depress Anxiety* 2009;26:888-901.

Assendelft WJ, Morton SC, Yu EI, et al. Spinal manipulative therapy for low back pain. *Cochrane Database Syst Rev* 2004;(1):CD000447.

[Return to Table of Contents](#)

www.icsi.org

References

- Atluri S, Akbik H, Sudarshan G. Prevention of opioid abuse in chronic non-cancer pain: an algorithmic, evidence based approach. *Pain Physician* 2012;15:ES177-89.
- Atluri S, Glaser SE, Shah RV, Sudarshan G. Needle position analysis in cases of paralysis from transforaminal epidurals: consider alternative approaches to traditional technique. *Pain Physician* 2013;16:321-34.
- Bair MJ, Robinson RL, Katon W, Kroenke K. Depression and pain comorbidity: a literature review. *Arch Intern Med* 2003;163:2433-45.
- Bakar Y, Sertel M, Oztürk A, et al. Short term effects of classic massage compared to connective tissue massage on pressure pain threshold and muscle relaxation response in women with chronic neck pain: a preliminary study. *J Manipulative Physiol Ther* 2014;37:415-21.
- Bartleson JD, Deen HG. *In Spine Disorders: Medical and Surgical Management*. 2009.
- Baskurt F, Ozcan A, Algun C. Comparison of effects of phonophoresis and iontophoresis of naproxen in the treatment of lateral epicondylitis. *Clin Rehabil* 2003;17:96-100.
- Bates C, Laciak R, Southwick A, Bishoff J. Overprescription of postoperative narcotics: a look at postoperative pain medication delivery, consumption and disposal in urological practice. *J Urol* 2011;185:551-55.
- Becker WC, Meghani SH, Barth KS, et al. Characteristics and outcomes of patients discharged from the opioid renewal clinic at the Philadelphia VA medical center. *Am J Addict* 2009;18:135-39.
- Benyamin R, Trescot AM, Datta S, et al. Opioid complications and side effects. *Pain Physician* 2008;11:S105-20.
- Benzon HT, Huntoon MA, Rathmell JP. Improving the safety of epidural steroid injections. *JAMA* 2015;313:1713-14.
- Berna C, Kulich RJ, Rathmell JP. Tapering long-term opioid therapy in chronic noncancer pain: evidence and recommendations for everyday practice. *Mayo Clin Proc* 2015;90:828-42.
- Berry CR, Chapman ED, Covington JL, et al. *In Pain: Current Understanding of Assessment, Management, and Treatments*. 2001.
- Bicket MC, Horowitz JM, Benzon HT, Cohen SP. Epidural injections in prevention of surgery for spinal pain: systematic review and meta-analysis of randomized controlled trials. *Spine J* 2015;15:348-62.
- Birse F, Derry S, Moore RA. Phenytoin for neuropathic pain and fibromyalgia in adults. *Cochrane Database Syst Rev* 2012;(5):CD009485.
- Bock E. Prescription opioid use may be decreasing, but heroin use is increasing. *NIH* 2015;25.
- Bohnert AS, Logan JE, Ganoczy D, Dowell D. A detailed exploration into the association of prescribed opioid dosage and overdose deaths among patients with chronic pain. *Med Care* 2016;54:435-41.
- Bohnert AS, Valenstein M, Bair MJ, et al. Association between opioid prescribing patterns and opioid overdose-related deaths. *JAMA* 2011;305:1315-21.
- Bolash R, Udeh B, Saweris Y, et al. Longevity and cost of implantable intrathecal drug delivery systems for chronic pain management: a retrospective analysis of 365 patients. *Neuromodulation* 2015;18:150-56.
- Borenstein DG, Lacks S, Wiesel SW. Cyclobenzaprine and naproxen versus naproxen alone in the treatment of acute low back pain and muscle spasm. *Clin Ther* 1990;12:125-31.
- Borman P, Keskin D, Ekici B, Bodur H. The efficacy of intermittent cervical traction in patents with chronic neck pain. *Clin Rheumatol* 2008;27:1249-53.

[Return to Table of Contents](#)

References

Boscarino JA, Rukstalis MR, Hoffman SN, et al. Prevalence of prescription opioid-use disorder among chronic pain patients: comparison of the DSM-5 vs. DSM-4 diagnostic criteria. *J Addict Dis* 2011;30:185-94.

Boyer EW. Management of opioid analgesic overdose. *N Engl J Med* 2012;367:146-55.

Brattberg G. Connective tissue massage in the treatment of fibromyalgia. *Eur J Pain* 1999;3:235-44.

Brecht S, Courtecuisse C, Debieuvre C, et al. Efficacy and safety of duloxetine 60 mg once daily in the treatment of pain in patients with major depressive disorder and at least moderate pain of unknown etiology: a randomized controlled trial. *J Clin Psychiatry* 2007;68:1707-16.

Breslau N, Davis GC, Andreski P. Migraine, psychiatric disorders, and suicide attempts: an epidemiologic study of young adults. *Psychiatry Res* 1991;37:11-23.

Briggs GG, Freeman RK. *In Drugs in Pregnancy and Lactation: a Reference Guide to Fetal and Neonatal Risk*. Tenth Edition. 2014.

Broderick JE, Junghaenel DU, Schwartz JE. Written emotional expression produces health benefits in fibromyalgia patients. *Psychosom Med* 2005;67:326-34.

Broussard CS, Rasmussen SA, Reefhuis J, et al. Maternal treatment with opioid analgesics and risk for birth defects. *Am J Obstet Gynecol* 2011;204:314.e1-11.

Büchi S, Buddeberg C, Klaghofer R, et al. Preliminary validation of PRISM (pictorial representation of illness and self measure) – a brief method to assess suffering. *Psychother Psychosom* 2002;71:333-41.

Buhrman M, Syk M, Burvall O, et al. Individualized guided internet-delivered cognitive-behavior therapy for chronic pain patients with comorbid depression and anxiety: a randomized controlled trial. *Clin J Pain* 2015;31:504-16.

Buppert C. New standard of care for prescribing opioids. 2015.

Butler M, Krebs E, Sunderlin B, Kane RL. Medical cannabis for non-cancer pain: a systematic review. 2015.

Butler SF, Cassidy TA, Chilcoat H, et al. Abuse rates and routes of administration of reformulated extended-release oxycodone: initial findings from a sentinel surveillance sample of individuals assessed for substance abuse treatment. *J Pain* 2013;14:351-58.

Butrans (package insert). LTS Lohmann Therapy Systems Corp., West Caldwell, NJ; 2014. Available at: <http://druginserts.com/lib/rx/meds/butrans-1/>.

Carrieri MP, Amass L, Lucas GM, et al. Buprenorphine use: the international experience. *Clin Infect Dis* 2006;43:S197-215.

Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep* 1985;100:126-31.

Cassidy TA, DasMahapatra P, Black RA, et al. Changes in prevalence of prescription opioid abuse after introduction of an abuse-deterrent opioid formulation. *Pain Med* 2014;15:440-51.

Castro MM, Daltro C, Kraychete DC, Lopes J. The cognitive behavioral therapy causes an improvement in quality-of-life in patients with chronic musculoskeletal pain. *Arq Neuropsiquiatr* 2012;70:864-68.

Centers for Disease Control and Prevention. Adult use of prescription opioid pain medications – Utah, 2008. *MMWR Morb Mortal Wkly Rep* 2010;59:153-57.

Centers for Disease Control and Prevention. CDC guideline for prescribing opioids for chronic pain – United States, 2016. *MMWR* 2016;65:1-49.

[Return to Table of Contents](#)

References

Centers for Disease Control and Prevention. Community-based opioid overdose prevention programs providing naloxone – United States, 2010. *MMWR Morb Mortal Wkly Rep* 2012a;61:101-05.

Centers for Disease Control and Prevention. Drug overdose deaths in the United States hit record numbers in 2014. 2015.

Centers for Disease Control and Prevention. Vital signs: overdoses of prescription opioid pain relievers – United States, 1999-2008. *MMWR Morb Mortal Wkly Rep* 2011;60:1487-92.

Chaparro LE, Furlan AD, Deshpande A, et al. Opioids compared with placebo or other treatments for chronic low back pain: an update of the Cochrane review. *Spine* 2014;39:556-63.

Cheatle MD, Savage SR. Informed consent in opioid therapy: a potential obligation and opportunity. *J Pain Symptom Manage* 2012;44:105-16.

Chen JH, Humphreys K, Shah NH, Lembke A. Distribution of opioids by different types of Medicare prescribers. *JAMA Intern Med* 2016;176:259-61.

Cherkin DC, Deyo RA, Battié M, et al. A comparison of physical therapy, chiropractic manipulation, and provision of an educational booklet for the treatment of patients with low back pain. *N Engl J Med* 1998;339:1021-29.

Cherkin DC, Eisenberg D, Sherman KJ, et al. Randomized trial comparing traditional Chinese medical acupuncture, therapeutic massage, and self-care education for chronic low back pain. *Arch Intern Med* 2001;161:1081-88.

Cherkin DC, Sherman KJ, Balderson BH, et al. Effect of mindfulness-based stress reduction vs cognitive behavioral therapy or usual care on back pain and functional limitations in adults with chronic low back pain: a randomized clinical trial. *JAMA* 2016;315:1240-49.

Chou R, Ballantyne JC, Fanciullo GJ, et al. Research gaps on use of opioids for chronic noncancer pain: findings from a review of the evidence for an American pain society and American academy of pain medicine clinical practice guideline. *J Pain* 2009a;10:147-59.

Chou R, Fanciullo GJ, Fine PG, et al. Clinical guidelines for the use of chronic opioid therapy in chronic noncancer pain. *J Pain* 2009b;10:113-30.

Chou R, Fanciullo GJ, Fine PG, et al. Opioids for chronic noncancer pain: prediction and identification of aberrant drug-related behaviors: a review of the evidence for an American pain society and American academy of pain medicine clinical practice guideline. *J Pain* 2009c;10:131-46.

Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American pain society. *Spine* 2009d;34:1066-77.

Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American college of physicians and the American pain society. *Ann Intern Med* 2007;147:478-91.

Chou R, Turner JA, Devine EB, et al. The effectiveness and risks of long-term opioid therapy for chronic pain: a systematic review for a national institutes of health pathways to prevention workshop. *Ann Intern Med* 2015;162:276-86.

Christie MJ. Cellular neuroadaptations to chronic opioids: tolerance, withdrawal and addiction. *Br J Pharmacol* 2008;154:384-96.

Chu LF, Clark DJ, Angst MS. Opioid tolerance and hyperalgesia in chronic pain patients after one month of oral morphine therapy: a preliminary prospective study. *J Pain* 2006;7:43-48.

Cicero TJ, Ellis MS, Surratt HL. Effect of abuse-deterrent formulation of oxycontin. *N Engl J Med* 2012;367:187-89.

[Return to Table of Contents](#)

References

- Cicero TJ, Ellis MS, Surratt HL, Kurtz SP. The changing face of heroin use in the United States: a retrospective analysis of the past 50 years. *JAMA Psychiatry* 2014;71:821-26.
- Cifu DX. In *Braddom's Physical Medicine and Rehabilitation, 5e*. 5th Edition. 2016.
- Cleary BJ, Donnelly JM, Strawbridge JD, et al. Methadone and perinatal outcomes: a retrospective cohort study. *Am J Obstet Gynecol* 2011;204:139.e1-9.
- Codeine (package insert). Roxane Laboratories, Inc., Columbus, OH; 2015. Available at: <http://druginserts.com/lib/rx/meds/codeine-sulfate-3/page/3/>.
- Coffin P, Banta-Green C. The dueling obligations of opioid stewardship. *Ann Intern Med* 2014;160:207.
- Coffin PO, Behar E, Rowe C, et al. Nonrandomized intervention study of naloxone coprescription for primary care patients receiving long-term opioid therapy for pain. *Ann Intern Med* 2016;165:245-52.
- Coffin PO, Sullivan SD. Cost-effectiveness of distributing naloxone to heroin users for lay overdose reversal. *Ann Intern Med* 2013;158:1-9.
- Cohen SP, Bicket MC, Jamison D, et al. Epidural steroids: a comprehensive, evidence-based review. *Reg Anesth Pain Med* 2013a;38:175-200.
- Cohen SP, Chen Y, Neufeld NJ. Sacroiliac joint pain: a comprehensive review of epidemiology, diagnosis and treatment. *Expert Rev Neurother* 2013b;13:99-116.
- Cohen SP, Stojanovic MP, Crooks M, et al. Lumbar zygapophysial (facet) joint radio frequency denervation success as a function of pain relief during diagnostic medial branch blocks: a multicenter analysis. *Spine J* 2008;8:498-504.
- Cohen SP, Wenzell D, Hurley RW, et al. A double-blind, placebo-controlled, dose-response pilot study evaluating intradiscal etanercept in patients with chronic discogenic low back pain or lumbosacral radiculopathy. *Anesthesiology* 2007;107:99-105.
- Cohen SP, White RL, Kurihara C, et al. Epidural steroids, etanercept, or saline in subacute sciatica: a multicenter, randomized trial. *Ann Intern Med* 2012;156:551-59.
- Cohen SP, Williams KA, Kurihara C, et al. Multicenter, randomized, comparative cost-effectiveness study comparing 0, 1, and 2 diagnostic medial branch (facet joint nerve) block treatment paradigms before lumbar facet radio frequency denervation. *Anesthesiology* 2010;113:395-405.
- Colebatch AN, Marks JL, van der Heijde DM, Edwards CJ. Safety of nonsteroidal antiinflammatory drugs and/or paracetamol in people receiving methotrexate for inflammatory arthritis: a Cochrane systematic review. *J Rheumatol Suppl* 2012;90:62-73.
- Compton WM, Jones CM, Baldwin GT. Relationship between nonmedical prescription-opioid use and heroin use. *N Engl J Med* 2016;374:154-63.
- Coplan PM, Kale H, Sandstrom L, et al. Changes in oxycodone and heroin exposures in the national poison data system after introduction of extended-release oxycodone with abuse-deterrent characteristics. *Pharmacoepidemiol Drug Saf* 2013;22:1274-82.
- Cousins G, Boland F, Courtney B, et al. Risk of mortality on and off methadone substitution treatment in primary care: a national cohort study. *Addiction* 2016;111:73-82.
- Cramer H, Lauche R, Hohmann C, et al. Randomized-controlled trial comparing yoga and home-based exercise for chronic neck pain. *Clin J Pain* 2013;29:216-23.
- Crews KR, Gaedigk A, Dunnenberger HM, et al. Clinical pharmacogenetics implementation consortium guidelines for cytochrome P450 2D6 genotype and codeine therapy: 2014 update. *Clin Pharmacol Ther* 2014;95:376-82.

References

Crider AB, Glaros AG. A meta-analysis of EMG biofeedback treatment of temporomandibular disorders. *J Orofac Pain* 1999;13:29-37.

Cuesta-Vargas AI, García-Romero JC, Arroyo-Morales M, et al. Exercise, manual therapy, and education with or without high-intensity deep-water running for non-specific chronic low back pain: a pragmatic randomized controlled trial. *Am J Phys Med Rehabil* 2011;90:526-34.

Cuesta-Vargas AI, White M, González-Sánchez M, Kuisma R. The optimal frequency of aquatic physiotherapy for individuals with chronic musculoskeletal pain: a randomised controlled trial. *Disabil Rehabil* 2015;37:311-18.

Cummings TM, White AR. Needling therapies in the management of myofascial trigger point pain: a systematic review. *Arch Phys Med Rehabil* 2001;82:986-92.

Dagenais S, Tricco AC, Haldeman S. Synthesis of recommendations for the assessment and management of low back pain from recent clinical practice guidelines. *Spine J* 2010;10:514-29.

Daniels SE, Riff D, Diamond E, et al. An assessment of the efficacy and safety of diclofenac potassium liquid-filled capsules in patients with various levels of baseline pain intensity. *Curr Med Res Opin* 2012;28:953-61.

Dart RC, Surratt HL, Cicero TJ, et al. Trends in opioid analgesic abuse and mortality in the United States. *N Engl J Med* 2015;372:241-48.

Dasgupta N, Funk MJ, Proescholdbell S, et al. Cohort study of the impact of high-dose opioid analgesics on overdose mortality. *Pain Med* 2016;17:85-98.

Daubresse M, Chang HY, Yu Y, et al. Ambulatory diagnosis and treatment of nonmalignant pain in the United States, 2000-2010. *Med Care* 2013;51:870-78.

de Rooij A, van der Leeden M, Roorda LD, et al. Predictors of outcome of multidisciplinary treatment in chronic widespread pain: an observational study. *BMC Musculoskelet Disord* 2013;14:133.

Dean M. Opioids in renal failure and dialysis patients. *J Pain Symptom Manage* 2004;28:497-504.

Deer TR, Mekhail N, Provenzano D, et al. The appropriate use of neurostimulation of the spinal cord and peripheral nervous system for the treatment of chronic pain and ischemic diseases: the neuro-modulation appropriateness consensus committee. *Neuromodulation* 2014;17:515-50.

Deer TR, Prager J, Levy R, et al. Polyanalgesic consensus conference 2012: recommendations for the management of pain by intrathecal (intraspinal) drug delivery: report of an interdisciplinary expert panel. *Neuromodulation* 2012;15:436-66.

DeNoon DJ. The 10 most prescribed drugs: most-prescribed drug list differs from list of drugs with biggest market share. 2011.

Derry S, Moore RA. Topical capsaicin (low concentration) for chronic neuropathic pain in adults. *Cochrane Database Syst Rev* 2012;(9):CD010111.

Desai MJ, Kapural L, Petersohn JD, et al. A prospective, randomized, multicenter, open-label clinical trial comparing intradiscal biacuplasty to conventional medical management for discogenic lumbar back pain. *Spine* 2016;41:1065-74.

Desai RJ, Hernandez-Diaz S, Bateman BT, Huybrechts KF. Increase in prescription opioid use during pregnancy among Medicaid-enrolled women. *Obstet Gynecol* 2014;123:997-1002.

Desai RJ, Huybrechts KF, Hernandez-Diaz S, et al. Exposure to prescription opioid analgesics in utero and risk of neonatal abstinence syndrome: population based cohort study. *BMJ* 2015;350:h2102.

[Return to Table of Contents](#)

References

- Devers A, Galer BS. Topical lidocaine patch relieves a variety of neuropathic pain conditions: an open-label study. *Clin J Pain* 2000;16:205-08.
- DeVries A, Koch T, Wall E, et al. Opioid use among adolescent patients treated for headache. *J Adolesc Health* 2014;55:128-33.
- Deyo RA, Mirza SK, Turner JA, Martin BI. Overtreating chronic back pain: time to back off? *J Am Board Fam Med* 2009;22:62-68.
- Dhalla IA, Mamdani MM, Sivilotti ML, et al. Prescribing of opioid analgesics and related mortality before and after the introduction of long-acting oxycodone. *CMAJ* 2009;181:891-96.
- Dhalla IA, Persaud N, Juurlink DN. Facing up to the prescription opioid crisis. *BMJ* 2011;343:d5142.
- Dietrich T, Leeson R, Gugliotta B, Petersen B. Efficacy and safety of low dose subcutaneous diclofenac in the management of acute pain: a randomized double-blind trial. *Pain Pract* 2014;14:315-23.
- Dilaudid (package insert). Hospira, Inc., Lake Forest, IL 60045; 2015. Available at: <http://druginserts.com/lib/rx/meds/dilaudid-3/>.
- Dillworth T, Mendoza ME, Jensen MP. Neurophysiology of pain and hypnosis for chronic pain. *Transl Behav Med* 2012;2:65-72.
- Dowell D, Kunins HV, Farley TA. Opioid analgesics – risky drugs, not risky patients. *JAMA* 2013;309:2219-20.
- Drug Enforcement Administration. DEA releases new rules that create convenient but safe and secure prescription drug disposal options. Available at: <https://www.dea.gov/divisions/hq/2014/hq090814.shtml>. 2014a.
- Drug Enforcement Administration. Tramadol (trade names: Ultram®, Ultracet®). 2014b.
- Dufour N, Thamsborg G, Oefeldt A, et al. Treatment of chronic low back pain: a randomized, clinical trial comparing group-based multidisciplinary biopsychosocial rehabilitation and intensive individual therapist-assisted back muscle strengthening exercises. *Spine* 2010;35:469-76.
- Dumas EO, Pollack GM. Opioid tolerance development: a pharmacokinetic/pharmacodynamic perspective. *AAPS J* 2008;10:537-51.
- Dundar U, Solak O, Yigit I, et al. Clinical effectiveness of aquatic exercise to treat chronic low back pain: a randomized controlled trial. *Spine* 2009;34:1436-40.
- Dunn KM, Saunders KW, Rutter CM, et al. Opioid prescriptions for chronic pain and overdose: a cohort study. *Ann Intern Med* 2010;152:85-92.
- DuPen A, Shen D, Ersek M. Mechanisms of opioid-induced tolerance and hyperalgesia. *Pain Manag Nurs* 2007;8:113-21.
- Duragesic (package insert). ALZA Corporation, Vacaville, CA; 2014. Available at: <http://druginserts.com/lib/rx/meds/duragesic/>.
- Eccleston C, Malleon PN, Clinch J, et al. Chronic pain in adolescents: evaluation of a programme of interdisciplinary cognitive behaviour therapy. *Arch Dis Child* 2003;88:881-85.
- Elkins G, Jensen MP, Patterson DR. Hypnotherapy for the management of chronic pain. *Int J Clin Exp Hypn* 2007;55:275-87.
- Ellickson PL, D'Amico EJ, Collins RL, Klein DJ. Marijuana use and later problems: when frequency of recent use explains age of initiation effects (and when it does not). *Subst Use Misuse* 2005;40:343-59.

[Return to Table of Contents](#)

References

Exalgo (package insert). Mallinckrodt Brand Pharmaceuticals, Inc., Hazelwood, MO; 2015. Available at: <http://druginserts.com/lib/rx/meds/exalgo-3/>.

Falla D, Lindstrøm R, Rechter L, et al. Effectiveness of an 8-week exercise programme on pain and specificity of neck muscle activity in patients with chronic neck pain: a randomized controlled study. *Eur J Pain* 2013;17:1517-28.

Feldman N, Chatton A, Khan R, et al. Alcohol-related brief intervention in patients treated for opiate or cocaine dependence: a randomized controlled study. *Subst Abuse Treat Prev Policy* 2011;6:22.

Field T, Hernandez-Reif M, Seligman S, et al. Juvenile rheumatoid arthritis: benefits from massage therapy. *J Pediatr Psychol* 1997;22:607-17.

Fillingim RB, Bruehl S, Dworkin RH, et al. The ACTTION-American pain society pain taxonomy (AAPT): an evidence-based and multidimensional approach to classifying chronic pain conditions. *J Pain* 2014;15:241-49.

Fine PG, Portenoy RK; Ad Hoc Expert Panel on Evidence Review and Guidelines for Opioid Rotation. Establishing "best practices" for opioid rotation: conclusions of an expert panel. *J Pain Symptom Manage* 2009;38:418-25.

Fleming MF, Davis J, Passik SD. Reported lifetime aberrant drug-taking behaviors are predictive of current substance use and mental health problems in primary care patients. *Pain Med* 2008;9:1098-1106.

Frampton M. Experience assessment and management of pain in people with dementia. *Age Ageing* 2003;32:248-51.

Franklin GM, Mai J, Turner J, et al. Bending the prescription opioid dosing and mortality curves: impact of the Washington state opioid dosing guideline. *Am J Ind Med* 2012;55:325-31.

Freedman BA, Cohen SP, Kuklo TR, et al. Intradiscal electrothermal therapy (IDET) for chronic low back pain in active-duty soldiers: 2-year follow-up. *Spine J* 2003;3:502-09.

Freeman BJ, Fraser RD, Cain CM, et al. A randomized, double-blind, controlled trial: intradiscal electrothermal therapy versus placebo for the treatment of chronic discogenic low back pain. *Spine* 2005;30:2369-78.

Freeman BJ, Ludbrook GL, Hall S, et al. Randomized, double-blind, placebo-controlled, trial of transforaminal epidural etanercept for the treatment of symptomatic lumbar disc herniation. *Spine* 2013;38:1986-94.

French DJ, Holroyd KA, Pinell C, et al. Perceived self-efficacy and headache-related disability. *Headache* 2000;40:647-56.

Frey ME, Manchikanti L, Benyamin RM, et al. Spinal cord stimulation for patients with failed back surgery syndrome: a systematic review. *Pain Physician* 2009;12:379-97.

Friedly JL, Comstock BA, Turner JA, et al. A randomized trial of epidural glucocorticoid injections for spinal stenosis. *N Engl J Med* 2014;371:11-21.

Friedman BW, Dym AA, Davitt M, et al. Naproxen with cyclobenzaprine, oxycodone/acetaminophen, or placebo for treating acute low back pain: a randomized clinical trial. *JAMA* 2015;314:1572-80.

Fullerton CA, Kim M, Thomas CP, et al. Medication-assisted treatment with methadone: assessing the evidence. *Psychiatr Serv* 2014;65:146-57.

Furlan AD, Yazdi F, Tsertsvadze A, et al. A systematic review and meta-analysis of efficacy, cost-effectiveness, and safety of selected complementary and alternative medicine for neck and low-back pain. *Evid Based Complement Alternat Med* 2012;2012:953139.

[Return to Table of Contents](#)

References

- Fusco BM, Giacobozzo M. Peppers and pain: the promise of capsaicin. *Drugs* 1997;53:909-14.
- Gagnier JJ, van Tulder MW, Berman B, Bombardier C. Herbal medicine for low back pain: a Cochrane review. *Spine* 2007;32:82-92.
- Gaither JR, Goulet JL, Becker WC, et al. The association between receipt of guideline-concordant long-term opioid therapy and all-cause mortality. *J Gen Intern Med* 2016;31:492-501.
- Gaskell H, Moore RA, Derry S, Stannard C. Oxycodone for neuropathic pain and fibromyalgia in adults. *Cochrane Database Syst Rev* 2014;(6):CD010692.
- Gatchel RJ, McGeary DD, McGeary CA, Lippe B. Interdisciplinary chronic pain management: past, present, and future. *Am Psychol* 2014;69:119-30.
- Gatchel RJ, Okifuji A. Evidence-based scientific data documenting the treatment and cost-effectiveness of comprehensive pain programs for chronic nonmalignant pain. *J Pain* 2006;7:779-93.
- Gatchel RJ, Peng YB, Peters ML, et al. The biopsychosocial approach to chronic pain: scientific advances and future directions. *Psychological Bulletin* 2007;133:581-624.
- Gaynor PJ, Gopal M, Zheng W, et al. A randomized placebo-controlled trial of duloxetine in patients with major depressive disorder and associated painful physical symptoms. *Curr Med Res Opin* 2011a;27:1849-58.
- Gaynor PJ, Gopal M, Zheng W, et al. Duloxetine versus placebo in the treatment of major depressive disorder and associated painful physical symptoms: a replication study. *Curr Med Res Opin* 2011b;27:1859-67.
- Gelot S, Nakhla E. Opioid dosing in renal and hepatic impairment. *US Pharm* 2014;39:34-38.
- Gendreau RM, Thorn MD, Gendreau JF, et al. Efficacy of milnacipran in patients with fibromyalgia. *J Rheumatol* 2005;32:1975-85.
- Gilbody S, Bower P, Fletcher J, et al. Collaborative care for depression: a cumulative meta-analysis and review of longer-term outcomes. *Arch Intern Med* 2006;166:2314-21.
- Gillis ME, Lumley MA, Mosley-Williams A, et al. The health effects of at-home written emotional disclosure in fibromyalgia: a randomized trial. *Ann Behav Med* 2006;32:135-46.
- Gomes T, Mamdani MM, Dhalla IA, et al. Opioid dose and drug-related mortality in patients with nonmalignant pain. *Arch Intern Med* 2011;171:686-91.
- Goodyear-Smith F, Halliwell J. Anticonvulsants for neuropathic pain: gaps in the evidence. *Clin J Pain* 2009;25:528-36.
- Gray SL, Anderson ML, Dublin S, et al. Cumulative use of strong anticholinergics and incident dementia: a prospective cohort study. *JAMA Intern Med* 2015;175:401-07.
- Grider JS, Manchikanti L, Carayannopoulos A, et al. Effectiveness of spinal cord stimulation in chronic spinal pain: a systematic review. *Pain Physician* 2016;19:E33-54.
- Griebeler ML, Morey-Vargas OL, Brito JP, et al. Pharmacologic interventions for painful diabetic neuropathy: an umbrella systematic review and comparative effectiveness network meta-analysis. *Ann Intern Med* 2014;161:639-49.
- Grossman P, Niemann L, Schmidt S, Walach H. Mindfulness-based stress reduction and health benefits: a meta-analysis. *J Psychosom Res* 2004;57:35-43.

[Return to Table of Contents](#)

References

- Grossman P, Tiefenthaler-Gilmer U, Raysz A, Kesper U. Mindfulness training as an intervention for fibromyalgia: evidence of postintervention and 3-year follow-up benefits in well-being. *Psychother Psychosom* 2007;76:226-33.
- Gudin JA. Clinical strategies for the primary health care professional to minimize prescription opioid abuse. *Postgrad Med* 2012;124:131-38.
- Gugelmann HM, Perrone J. Can prescription drug monitoring programs help limit opioid abuse? *JAMA* 2011;306:2258-59.
- Guillemette S, Witzke S, Leier J, et al. Medical cost impact of intrathecal drug delivery for noncancer pain. *Pain Med* 2013;14:504-15.
- Gupta G, Radhakrishna M, Chankowsky J, Asenjo JF. Methylene blue in the treatment of discogenic low back pain. *Pain Physician* 2012;15:333-38.
- Gurley BJ, Gardner SF, Hubbard MA. Content versus label claims in ephedra-containing dietary supplements. *Am J Health Syst Pharm* 2000;57:963-69.
- Gwira Baumblatt JA, Wiedeman C, Dunn JR, et al. High-risk use by patients prescribed opioids for pain and its role in overdose deaths. *JAMA Intern Med* 2014;174:796-801.
- Haddock CK, Rowan AB, Andrasik F, et al. Home-based behavioral treatments for chronic benign headache: a meta-analysis of controlled trials. *Cephalalgia* 1997;17:113-18.
- Haffey F, Brady RR, Maxwell S. A comparison of the reliability of smartphone apps for opioid conversion. *Drug Saf* 2013;36:111-17.
- Hall J, Swinkels A, Briddon J, McCabe CS. Does aquatic exercise relieve pain in adults with neurologic or musculoskeletal disease? A systematic review and meta-analysis of randomized controlled trials. *Arch Phys Med Rehabil* 2008;89:873-83.
- Han B, Compton WM, Jones CM, Cai R. Nonmedical prescription opioid use and use disorders among adults aged 18 through 64 years in the United States, 2003-2013. *JAMA* 2015;314:1468-78.
- Hansen H, Manchikanti L, Simopoulos TT, et al. A systematic evaluation of the therapeutic effectiveness of sacroiliac joint interventions. *Pain Physician* 2012;15:E247-78.
- Hasenbring M, Hallner D, Klasen B. Psychological mechanisms in the transition from acute to chronic pain: over- or underrated? *Schmerz* 2001;15:442-47.
- Hatheway JA, Caraway D, David G, et al. Systemic opioid elimination after implantation of an intrathecal drug delivery system significantly reduced health-care expenditures. *Neuromodulation* 2015;18:207-13.
- Havens JR, Leukefeld CG, DeVaugh-Geiss AM, et al. The impact of a reformulation of extended-release oxycodone designed to deter abuse in a sample of prescription opioid abusers. *Drug Alcohol Depend* 2014;139:9-17.
- Hawkins RMF. A systematic meta-review of hypnosis as an empirically supported treatment for pain. *Pain Reviews* 2001;8:47-73.
- Hayden JA, Chou R, Hogg-Johnson S, Bombardier C. Systematic reviews of low back pain prognosis had variable methods and results: guidance for future prognosis reviews. *J Clin Epidemiol* 2009;62:781-796.
- Hayden JA, van Tulder MW, Malmivaara A, Koes BW. Exercise therapy for treatment of non-specific low back pain. *Cochrane Database Syst Rev* 2011;(3):CD000335.
- Hayes SC, Strosahl KD, Wilson K. *In Acceptance and Commitment Therapy: An Experiential Approach to Behavior Change*. New York, NY, US: Guilford Press. 1999.

[Return to Table of Contents](#)

References

- Hearn L, Derry S, Moore RA. Lacosamide for neuropathic pain and fibromyalgia in adults. *Cochrane Database Syst Rev* 2012;(2):CD009318.
- Hegmann KT, Weiss MS, Bowden K, et al. ACOEM practice guidelines: opioids for treatment of acute, subacute, chronic, and postoperative pain. *J Occup Environ Med* 2014;56:e143-59.
- Heit HA, Gourlay DL. Urine drug testing in pain medicine. *J Pain Symptom Manage* 2004;27:260-67.
- Henriques AA, Dussán-Sarria JA, Botelho LM, Caumo W. Multidimensional approach to classifying chronic pain conditions – less is more. *J Pain* 2014;15:1199-200.
- Heymans MW, van Buuren S, Knol DL, et al. The prognosis of chronic low back pain is determined by changes in pain and disability in the initial period. *Spine J* 2010;10:847-56.
- Hochberg MC, Altman RD, April KT, et al. American college of rheumatology 2012 recommendations for the use of nonpharmacologic and pharmacologic therapies in osteoarthritis of the hand, hip, and knee. *Arthritis Care Res* 2012;64:465-74.
- Hoge CW, Terhakopian A, Castro CA, et al. Association of posttraumatic stress disorder with somatic symptoms, health care visits, and absenteeism among Iraq war veterans. *Am J Psychiatry* 2007;164:150-53.
- Holbech JV, Otto M, Bach FW, et al. The anticonvulsant levetiracetam for the treatment of pain in polyneuropathy: a randomized, placebo-controlled, cross-over trial. *Eur J Pain* 2011;15:608-14.
- Holmberg C, Rappenecker J, Karner JJ, Witt CM. The perspectives of older women with chronic neck pain on perceived effects of qigong and exercise therapy on aging: a qualitative interview study. *Clin Interv Aging* 2014;9:403-10.
- Hooten WM. Chronic pain and mental health disorders: shared neural mechanisms, epidemiology, and treatment. *Mayo Clin Proc* 2016;91:955-70.
- Hooten WM, Lamer TJ, Twyner C. Opioid-induced hyperalgesia in community-dwelling adults with chronic pain. *Pain* 2015a;156:1145-52.
- Hooten WM, St. Sauver JL, McGree ME, et al. Incidence and risk factors for progression from short-term to episodic or long-term opioid prescribing: a population-based study. *Mayo Clin Proc* 2015b;90:850-56.
- Hopwood MB, Abram SE. Factors associated with failure of lumbar epidural steroids. *Reg Anesth* 1993;18:238-43.
- Hsieh LL, Kuo CH, Lee LH, et al. Treatment of low back pain by acupuncture and physical therapy: randomised controlled trial. *BMJ* 2006;332:696-700.
- Hurwitz EL, Carragee EJ, van der Velde G, et al. Treatment of neck pain: noninvasive interventions: results of the bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *Spine* 2008;33:S123-52.
- Hwang CS, Chang HY, Alexander GC. Impact of abuse-deterrent OxyContin on prescription opioid utilization. *Pharmacoepidemiol Drug Saf* 2015;24:197-204.
- Hyslinga (package insert). Purdue Pharma L.P., Stamford, CT; 2014. Available at: <http://app.purduepharma.com/xmlpublishing/pi.aspx?id=h>.
- International Association for the Study of Pain. Classification of chronic pain: descriptions of chronic pain syndromes and definitions of pain terms. Second Edition. Available at: <http://www.iasp-pain.org/PublicationsNews/Content.aspx?ItemNumber=1673>. Updated August 6, 2014.
- Ives TJ, Chelminski PR, Hammett-Stabler CA, et al. Predictors of opioid misuse in patients with chronic pain: a prospective cohort study. *BMC Health Serv Res* 2006;6:46.

References

Jamison RN, VadeBoncouer T, Ferrante FM. Low back pain patients unresponsive to an epidural steroid injection: identifying predictive factors. *Clin J Pain* 1991;7:311-17.

Janssens KAM, Zijlema WL, Joustra ML, Rosmalen JGM. Mood and anxiety disorders in chronic fatigue syndrome, fibromyalgia, and irritable bowel syndrome: results from the lifelines cohort study. *Psychosom Med* 2015;77:449-57.

Jensen MP. The neurophysiology of pain perception and hypnotic analgesia: implications for clinical practice. *Am J Clin Hypn* 2008;51:123-48.

Jensen MP, Patterson DR. Hypnotic approaches for chronic pain management: clinical implications of recent research findings. *Am Psychol* 2014;69:167-77.

Jensen MP, Turner JA, Romano JM, Karoly P. Coping with chronic pain: a critical review of the literature. *Pain* 1991;47:249-83.

Johnson H, Paulozzi L, Porucznik C, et al. Decline in drug overdose deaths after state policy changes – Florida, 2010-2012. *MMWR Morb Mortal Wkly Rep* 2014;63:569-74.

Johnson SJ. Opioid safety in patients with renal or hepatic dysfunction. Available at: <http://www.pain-topics.org>. 2007.

Jones CM. Heroin use and heroin use risk behaviors among nonmedical users of prescription opioid pain relievers – United States, 2002-2004 and 2008-2010. *Drug Alcohol Depend* 2013a;132:95-100.

Jones CM, Mack KA, Paulozzi LJ. Pharmaceutical overdose deaths, United States, 2010. *JAMA* 2013b;309:657-59.

Jones CM, McAninch JK. Emergency department visits and overdose deaths from combined use of opioids and benzodiazepines. *Am J Prev Med* 2015;49:493-501.

Jones T, Lookatch S, Grant P, et al. Further validation of an opioid risk assessment tool: the brief risk interview. *J Opioid Manag* 2014;10:353-64.

Jones T, Moore T, Levy JL, et al. A comparison of various risk screening methods in predicting discharge from opioid treatment. *Clin J Pain* 2012;28:93-100.

Jordan JL, Holden MA, Mason EE, Foster NE. Interventions to improve adherence to exercise for chronic musculoskeletal pain in adults. *Cochrane Database Syst Rev* 2010;(1):CD005956.

Jull GA, Falla D, Vicenzino B, Hodges PW. The effect of therapeutic exercise on activation of the deep cervical flexor muscles in people with chronic neck pain. *Man Ther* 2009;14:696-701.

Jüni P, Battaglia M, Nüesch E, et al. A randomised controlled trial of spinal manipulative therapy in acute low back pain. *Ann Rheum Dis* 2009;68:1420-27.

Juurlink DN, Dhalla IA. Dependence and addiction during chronic opioid therapy. *J Med Toxicol* 2012;8:393-99.

Kabat-Zinn J. *In Full Catastrophe Living: Using the Wisdom of Your Body and Mind to Face Stress, Pain, and Illness*. 2009.

Kalira V, Treisman GJ, Clark MR. Borderline personality disorder and chronic pain: a practical approach to evaluation and treatment. *Curr Pain Headache Rep* 2013;17:350.

Kamper SJ, Apeldoorn AT, Chiarotto A, et al. Multidisciplinary biopsychosocial rehabilitation for chronic low back pain. *Cochrane Database Syst Rev* 2014;(9):CD000963.

Kapural L, Vrooman B, Sarwar S, et al. Radio frequency intradiscal biacuplasty for treatment of discogenic lower back pain: a 12-month follow-up. *Pain Med* 2015a;16:425-31.

[Return to Table of Contents](#)

References

Kapural L, Yu C, Doust MW, et al. Novel 10-kHz high-frequency therapy (HF10 therapy) is superior to traditional low-frequency spinal cord stimulation for the treatment of chronic back and leg pain: the SENZA-RCT randomized controlled trial. *Anesthesiology* 2015b;123:851-60.

Katon W, Von Korff M, Lin E, et al. Stepped collaborative care for primary care patients with persistent symptoms of depression: a randomized trial. *Arch Gen Psychiatry* 1999;56:1109-15.

Keller S, Bann CM, Dodd SL, et al. Validity of the brief pain inventory for use in documenting the outcomes of patients with noncancer pain. *Clin J Pain* 2004;20:309-18.

Kennedy DJ, Engel A, Kreiner DS, et al. Fluoroscopically guided diagnostic and therapeutic intra-articular sacroiliac joint injections: a systematic review. *Pain Med* 2015;16:1500-18.

Khot A, Bowditch M, Powell J, Sharp D. The use of intradiscal steroid therapy for lumbar spinal discogenic pain: a randomized controlled trial. *Spine* 2004;29:833-37.

Kietrys DM, Palombaro KM, Azzaretto E, et al. Effectiveness of dry needling for upper-quarter myofascial pain: a systematic review and meta-analysis. *J Orthop Sports Phys Ther* 2013;43:620-34.

Kim SH, Ahn SH, Cho YW, Lee DG. Effect of intradiscal methylene blue injection for the chronic discogenic low back pain: one year prospective follow-up study. *Ann Rehabil Med* 2012;36:657-64.

Kirpalani D, Mitra R. Is chronic opioid use a negative predictive factor for response to cervical epidural steroid injections? *J Back Musculoskelet Rehabil* 2011;24:123-27.

Kirschner N, Ginsburg J, Sulmasy LS. Prescription drug abuse: executive summary of a policy position paper from the American college of physicians. *Ann Intern Med* 2014;160:198-200.

Knotkova H, Fine PG, Portenoy RK. Opioid rotation: the science and the limitations of the equianalgesic dose table. *J Pain Symptom Manage* 2009;38:426-39.

Koumantakis GA, Watson PJ, Oldham JA. Trunk muscle stabilization training plus general exercise versus general exercise only: randomized controlled trial of patients with recurrent low back pain. *Phys Ther* 2005;85:209-25.

Krantz MJ, Martin J, Stimmel B, et al. QTc interval screening in methadone treatment. *Ann Intern Med* 2009;150:387-95.

Krebs EE, Lorenz KA, Bair MJ, et al. Development and initial validation of the PEG, a three-item scale assessing pain intensity and interference. *J Gen Intern Med* 2009;24:733-38.

Kroenke K, Bair MJ, Damush TM, et al. Optimized antidepressant therapy and pain self-management in primary care patients with depression and musculoskeletal pain: a randomized controlled trial. *JAMA* 2009a;301:2099-110.

Kroenke K, Bair M, Damush T, et al. Stepped care for affective disorders and musculoskeletal pain (SCAMP) study: design and practical implications of an intervention for comorbid pain and depression. *Gen Hosp Psychiatry* 2007;29:506-17.

Kroenke K, Krebs EE, Bair MJ. Pharmacotherapy of chronic pain: a synthesis of recommendations from systematic reviews. *Gen Hosp Psychiatry* 2009b;31:206-19.

Kroenke K, Shen J, Oxman TE, et al. Impact of pain on the outcomes of depression treatment: results from the RESPECT trial. *Pain* 2008;134:209-15.

Kronick R. AHRQ data reveal wider impact of opioid overuse. Available at: <http://www.ahrq.gov/news/blog/ahrqviews/100914.html>. 2014.

Krupitsky E, Nunes EV, Ling W, et al. Injectable extended-release naltrexone (XR-NTX) for opioid dependence: long-term safety and effectiveness. *Addiction* 2013;108:1628-37.

References

- Kuehn BM. CDC: Major disparities in opioid prescribing among states: some states crack down on excess prescribing. *JAMA* 2014;312:684-86.
- Kuehn BM. SAMHSA: Pain medication abuse a common path to heroin: experts say this pattern likely driving heroin resurgence. *JAMA* 2013;310:1433-34.
- Kvarstein G, Måwe L, Indahl A, et al. A randomized double-blind controlled trial of intra-annular radio frequency thermal disc therapy – a 12-month follow-up. *Pain* 2009;145:279-86.
- Lam M, Galvin R, Curry P. Effectiveness of acupuncture for non-specific chronic low back pain: a systematic review and meta-analysis. *Spine* 2013;38:2124-38.
- Lanier WL, Kharasch ED. Contemporary clinical opioid use: opportunities and challenges. *Mayo Clin Proc* 2009;84:572-75.
- Larochelle MR, Liebschutz JM, Zhang F, et al. Opioid prescribing after non-fatal overdose and association with repeated overdose: a cohort study. *Ann Intern Med* 2016;164:1-9.
- Lee JH, Choi TY, Lee MS, et al. Acupuncture for acute low back pain: a systematic review. *Clin J Pain* 2013;29:172-85.
- Lembke A. Why doctors prescribe opioids to known opioid abusers. *N Engl J Med* 2012 ;367:1580-81.
- Lépine JP, Briley M. The epidemiology of pain in depression. *Hum Psychopharmacol* 2004;19:S3-7.
- Leppin AL, Montori VM, Gionfriddo MR. Minimally disruptive medicine: a pragmatically comprehensive model for delivering care to patients with multiple chronic conditions. *Healthcare* 2015;3:50-63.
- Lewis NL, Williams JE. Acute pain management in patients receiving opioids for chronic and cancer pain. *Cont Educ in Anaesthesia, Crit Care & Pain* 2005;5:127-29.
- Liang Y, Turner BJ. Assessing risk for drug overdose in a national cohort: role for both daily and total opioid dose? *J Pain* 2015;16:318-25.
- Liebschutz JM, Saitz R, Weiss RD, et al. Clinical factors associated with prescription drug use disorder in urban primary care patients with chronic pain. *J Pain* 2010;11:1047-55.
- Lindström I, Ohlund C, Eek C, et al. The effect of graded activity on patients with subacute low back pain: a randomized prospective clinical study with an operant-conditioning behavioral approach. *Phys Ther* 1992;72:279-90.
- Liu L, Huang QM, Liu QG, et al. Effectiveness of dry needling for myofascial trigger points associated with neck and shoulder pain: a systematic review and meta-analysis. *Arch Phys Med Rehabil* 2015;96:944-55.
- Liu W, Zahner L, Cornell M, et al. Benefit of Qigong exercise in patients with fibromyalgia: a pilot study. *Int J Neurosci* 2012;122:657-64.
- Liu X, Ye W, Watson P, Tepper P. Use of benzodiazepines, hypnotics, and anxiolytics in major depressive disorder: association with chronic pain diseases. *J Nerv Ment Dis* 2010;198:544-50.
- Loder E, Weizenbaum E, Frishberg B, Silberstein S. Choosing wisely in headache medicine: the American headache society's list of five things physicians and patients should question. *Headache* 2013;53:1651-59.
- Maas ET, Ostelo RW, Niemisto L, et al. Radio frequency denervation for chronic low back pain. *Cochrane Database Syst Rev* 2015;(10):CD008572.
- Maeda A, Bateman BT, Clancy CR, et al. Opioid abuse and dependence during pregnancy: temporal trends and obstetrical outcomes. *Anesthesiology* 2014;121:1158-65.

[Return to Table of Contents](#)

References

Magni G, Rigatti-Luchini S, Fracca F, Merskey H. Suicidality in chronic abdominal pain: an analysis of the Hispanic health and nutrition examination survey (HHANES). *Pain* 1998;76:137-44.

Makris UE, Abrams RC, Gurland B, Reid MC. Management of persistent pain in the older patient: a clinical review. *JAMA* 2014;312:825-36.

Malanga GA, Cruz Colon EJ. Myofascial low back pain: a review. *Phys Med Rehabil Clin N Am* 2010;21:711-24.

Malmivaara A, Koes BW, Bouter LM, van Tulder MW. Applicability and clinical relevance of results in randomized controlled trials: the Cochrane review on exercise therapy for low back pain as an example. *Spine* 2006;31:1405-09.

Manchikanti L, Abdi S, Atluri S, et al. American society of interventional pain physicians (ASIPP) guidelines for responsible opioid prescribing in chronic non-cancer pain: part I – evidence assessment. *Pain Physician* 2012a;15:S1-65.

Manchikanti L, Cash KA, Damron KS, et al. Controlled substance abuse and illicit drug use in chronic pain patients: an evaluation of multiple variables. *Pain Physician* 2006;9:215-25.

Manchikanti L, Falco FJ, Singh V, et al. Utilization of interventional techniques in managing chronic pain in the Medicare population: analysis of growth patterns from 2000 to 2011. *Pain Physician* 2012b;15:E969-82.

Manchikanti L, Kaye AD, Boswell MV, et al. A systematic review and best evidence synthesis of the effectiveness of therapeutic facet joint interventions in managing chronic spinal pain. *Pain Physician* 2015a;18:E535-82.

Manchikanti L, Nampiaparampil DE, Manchikanti KN, et al. Comparison of the efficacy of saline, local anesthetics, and steroids in epidural and facet joint injections for the management of spinal pain: a systematic review of randomized controlled trials. *Surg Neurol Int* 2015b;6:S194-235.

Mannion AF, Müntener M, Taimela S, Dvorak J. A randomized clinical trial of three active therapies for chronic low back pain. *Spine* 1999;24:2435-48.

Mao J. Clinical diagnosis of opioid-induced hyperalgesia. *Reg Anesth Pain Med* 2015;40:663-64.

Margetic P, Pavic R, Stancic MF. Provocative discography screening improves surgical outcome. *Wien Klin Wochenschr* 2013;125:600-10.

Martell BA, O'Connor PG, Kerns RD, et al. Systematic review: opioid treatment for chronic back pain: prevalence, efficacy, and association with addiction. *Ann Intern Med* 2007;146:116-27.

Mason L, Moore RA, Derry S, et al. Systematic review of topical capsaicin for the treatment of chronic pain. *BMJ* 2004;328:991.

McAlindon TE, Bannuru RR, Sullivan MC, et al. OARSI guidelines for the non-surgical management of knee osteoarthritis. *Osteoarthritis Cartilage* 2014;22:363-88.

McAuliffe L, Brown D, Fetherstonhaugh D. Pain and dementia: an overview of the literature. *Int J Older People Nurs* 2012;7:219-26.

McCarthy DM, Wolf MS, McConnell R, et al. Improving patient knowledge and safe use of opioids: a randomized controlled trial. *Acad Emerg Med* 2015;22:331-39.

McCleane GJ. Lamotrigine in the management of neuropathic pain: a review of the literature. *Clin J Pain* 2000;16:321-26.

McCracken LM. Learning to live with the pain: acceptance of pain predicts adjustment in persons with chronic pain. *Pain* 1998;74:21-27.

References

- McFarlane AC. The long-term costs of traumatic stress: intertwined physical and psychological consequences. *World Psychiatry* 2010;9:3-10.
- McFarlane AC, Atchison M, Rafalowicz E, Papay P. Physical symptoms in post-traumatic stress disorder. *J Psychosom Res* 1994;38:715-26.
- Medical Letter, The. Abuse-deterrent opioid formulations. *Med Lett Drugs Ther* 2015;57:119-21.
- Medical Letter, The. Naloxone (Narcan) nasal spray for opioid overdose. *Med Lett Drugs Ther* 2016;58:1-2.
- Meldrum ML. *In Opioids and Pain Relief: A Historical Perspective (Progress in Pain Research and Management, V. 25)*. 1st Edition. 2003.
- Meperidine (package insert). Epic Pharma, LLC Laurelton, NY; 2016. Available at: <http://druginserts.com/lib/rx/meds/meperidine-hydrochloride-5/>.
- Methadone (package insert). Major® Pharmaceuticals, Livonia, MI; 2016. Available at: <http://druginserts.com/lib/rx/meds/methadone-hydrochloride/>.
- Michna E, Jamison RN, Pham LD, et al. Urine toxicology screening among chronic pain patients on opioid therapy: frequency and predictability of abnormal findings. *Clin J Pain* 2007;23:173-79.
- Micromedex Solutions. Available at: <http://www.micromedex.com>. 2016.
- Miech R, Johnston L, O'Malley PM, et al. Prescription opioids in adolescence and future opioid misuse. *Pediatrics* 2015;136:e1169-77.
- Miller LG. Herbal medicinals: selected clinical considerations focusing on known or potential drug-herb interactions. *Arch Intern Med* 1998;158:2200-11.
- Miller M, Barber CW, Leatherman S, et al. Prescription opioid duration of action and the risk of unintentional overdose among patients receiving opioid therapy. *JAMA Intern Med* 2015;175:608-15.
- Minnesota Dental Association. MDA protocol for assessment and treatment of oral/facial pain. 2015.
- Moeller-Bertram T, Strigo IA, Simmons AN, et al. Evidence for acute central sensitization to prolonged experimental pain in posttraumatic stress disorder. *Pain Med* 2014;15:762-71.
- Moore PA, Hersh EV. Combining ibuprofen and acetaminophen for acute pain management after third-molar extractions: translating clinical research to dental practice. *J Am Dent Assoc* 2013;144:898-908.
- Moore TM, Jones T, Browder JH, et al. A comparison of common screening methods for predicting aberrant drug-related behavior among patients receiving opioids for chronic pain management. *Pain Med* 2009;10:1426-33.
- Moore RJ. *In Handbook of Pain and Palliative Care: Biobehavioral Approaches for the Life Course*. 2011.
- Morphine (package insert). Physicians Total Care, Inc. Tulsa, OK; 2012. Available at: <http://druginserts.com/lib/rx/meds/morphine-sulfate-24/>.
- Murray CJ, Atkinson C, Bhalla K, et al. The state of US health, 1990-2010: burden of diseases, injuries, and risk factors. *JAMA* 2013;310:591-608.
- Nagar VR, Springer JE, Salles S. Increased incidence of spinal abscess and substance abuse after implementation of state mandated prescription drug legislation. *Pain Med* 2015;16:2031-35.
- Nahin RL. Estimates of pain prevalence and severity in adults: United States, 2012. *J Pain* 2015;16:769-80.

[Return to Table of Contents](#)

References

- National Highway Traffic Safety Administration. Morphine (and heroin). Available at: <http://www.nhtsa.gov/people/injury/research/job185drugs/morphine.htm>. Accessed June 2016.
- Neumann AM, Blondell RD, Jaanimägi U, et al. A preliminary study comparing methadone and buprenorphine in patients with chronic pain and coexistent opioid addiction. *J Addict Dis* 2013;32:68-78.
- Nicholas MK, Asghari A, Corbett M, et al. Is adherence to pain self-management strategies associated with improved pain, depression and disability in those with disabling chronic pain? *Eur J Pain* 2012;16:93-104.
- Nielsen S, Degenhardt L, Hoban B, Gisev N. A synthesis of oral morphine equivalents (OME) for opioid utilisation studies. *Pharmacoepidemiol Drug Saf* 2016;25:733-37.
- Nirschl RP, Rodin DM, Ochiai DH, et al. Iontophoretic administration of dexamethasone sodium phosphate for acute epicondylitis. A randomized, double-blinded, placebo-controlled study. *Am J Sports Med* 2003;31:189-95.
- Noble M, Treadwell JR, Tregear SJ, et al. Long-term opioid management for chronic noncancer pain. *Cochrane Database Syst Rev* 2010;(1):CD006605.
- Norco (package insert). Actavis Laboratories FL, Inc. Fort Lauderdale, FL; 2014. Available at: <http://druginserts.com/lib/rx/meds/norco-2/>.
- Nuckols TK, Anderson L, Popescu I, et al. Opioid prescribing: a systematic review and critical appraisal of guidelines for chronic pain. *Ann Intern Med* 2014;160:38-47.
- Nucynta (package insert). Janssen Ortho, LLC, Gurabo, PR;2013. Available at: <http://druginserts.com/lib/rx/meds/nucynta-2/>.
- Nucynta ER (package insert). Janssen Ortho, LLC, Gurabo, PR;2014. Available at: <http://druginserts.com/lib/rx/meds/nucynta-er-1/>.
- Nygaard HA, Jarland M. Are nursing home patients with dementia diagnosis at increased risk for inadequate pain treatment? *Int J Geriatr Psychiatry* 2005;20:730-37.
- Ohtori S, Miyagi M, Eguchi Y, et al. Epidural administration of spinal nerves with the tumor necrosis factor-alpha inhibitor, etanercept, compared with dexamethasone for treatment of sciatica in patients with lumbar spinal stenosis: a prospective randomized study. *Spine* 2012;37:439-44.
- Okie S. A flood of opioids, a rising tide of deaths. *N Engl J Med* 2010;363:1981-85.
- Oltean H, Robbins C, van Tulder MW, et al. Herbal medicine for low-back pain. *Cochrane Database Syst Rev* 2014;(12):CD004504.
- Ong J, Claydon LS. The effect of dry needling for myofascial trigger points in the neck and shoulders: a systematic review and meta-analysis. *J Bodyw Mov Ther* 2014;18:390-98.
- Opana ER (package insert). Pharmaceuticals Manufacturing Research Services Inc., Horsham, PA; 2012. Available at: <http://druginserts.com/lib/rx/meds/opana-er-3/>.
- Oregon Health Authority. Oregon prescription drug monitoring program: 2012 annual report to the PDMP advisory commission. 2013.
- [Oslund S, Robinson RC, Clark TC, et al.](#) Long-term effectiveness of a comprehensive pain management program: strengthening the case for interdisciplinary care. *Proc (Bayl Univ MedCent)* 2009;22:211-14.
- Oxycodone (package insert). Mayne Pharma Montgomery, AL; 2014. Available at: <http://druginserts.com/lib/rx/meds/oxycodone-hydrochloride-10/>.

[Return to Table of Contents](#)

References

- Oxymorphone (package insert). CorePharma, LLC Middlesex, NJ; 2014. Available at: <http://druginserts.com/lib/rx/meds/oxymorphone-hydrochloride-4/>.
- Paras ML, Murad MH, Chen LP, et al. Sexual abuse and lifetime diagnosis of somatic disorders: a systematic review and meta-analysis. *JAMA* 2009;302:550-61.
- Park EM, Meltzer-Brody S, Suzuki J. Evaluation and management of opioid dependence in pregnancy. *Psychosomatics* 2012;53:424-32.
- Passik SD, Kirsh KL, Casper D. Addiction-related assessment tools and pain management: instruments for screening, treatment planning, and monitoring compliance. *Pain Med* 2008;9:S145-66.
- Pasternak GW. Molecular biology of opioid analgesia. *J Pain Symptom Manage* 2005;29:S2-9.
- Pasternak GW. Opiate pharmacology and relief of pain. *J Clin Oncol* 2014;32:1655-61.
- Patel VB, Manchikanti L, Singh V, et al. Systematic review of intrathecal infusion systems for long-term management of chronic non-cancer pain. *Pain Physician* 2009;12:345-60.
- Pathan H, Williams J. Basic opioid pharmacology: an update. *Br J Pain* 2012;6:11-16.
- Paulozzi LJ, Kilbourne EM, Desai HA. Prescription drug monitoring programs and death rates from drug overdose. *Pain Med* 2011;12:747-54.
- Paulozzi LJ, Kilbourne EM, Shah NG, et al. A history of being prescribed controlled substances and risk of drug overdose death. *Pain Med* 2012;13:87-95.
- Pauza KJ, Howell S, Dreyfuss P, et al. A randomized, placebo-controlled trial of intradiscal electrothermal therapy for the treatment of discogenic low back pain. *Spine J* 2004;4:27-35.
- Peng B, Pang X, Wu Y, et al. A randomized placebo-controlled trial of intradiscal methylene blue injection for the treatment of chronic discogenic low back pain. *Pain* 2010;149:124-29.
- Perahia DG, Quail D, Desai D, et al. Switching to duloxetine in selective serotonin reuptake inhibitor non- and partial-responders: effects on painful physical symptoms of depression. *J Psychiatr Res* 2009;43:512-18.
- Perlman AI, Sabina A, Williams AL, et al. Massage therapy for osteoarthritis of the knee: a randomized controlled trial. *Arch Intern Med* 2006;166:2533-38.
- Pesce A, West C, Rosenthal M, et al. Marijuana correlates with use of other illicit drugs in a pain patient population. *Pain Physician* 2010;13:283-87.
- Petrov ME, Sawyer P, Kennedy R, et al. Benzodiazepine (BZD) use in community-dwelling older adults: Longitudinal associations with mobility, functioning, and pain. *Arch Gerontol Geriatr* 2014;59:331-37.
- Pharmacist's Letter/Prescriber's Letter. Equianalgesic dosing of opioids for pain management. 2014.
- Pincus D, Sheikh AA. *In Imagery for Pain Relief: A Scientifically Grounded Guidebook for Clinicians*. Routledge. Taylor & Francis Group. New York, London. 2009.
- Pincus T, Burton AK, Vogel S, Field AP. A systematic review of psychological factors as predictors of chronicity/disability in prospective cohorts of low back pain. *Spine* 2002;27:E109-20.
- Pittler MH, Ernst E. Complementary therapies for neuropathic and neuralgic pain: systematic review. *Clin J Pain* 2008;24:731-33.
- Pope JE, Falowski S, Deer TR. Advanced waveforms and frequency with spinal cord stimulation: burst and high-frequency energy delivery. *Expert Rev Med Devices* 2015;12:431-37.
- Porter J, Jick H. Addiction rare in patients treated with narcotics. *N Engl J Med* 1980;302:123.

[Return to Table of Contents](#)

References

Prager J, Deer T, Levy R, et al. Best practices for intrathecal drug delivery for pain. *Neuromodulation* 2014;17:354-72.

Provenzano DA. Diagnostic discography: what is the clinical utility? *Curr Pain Headache Rep* 2012;16:26-34.

Rainville J, Jouve CA, Hartigan C, et al. Comparison of short- and long-term outcomes for aggressive spine rehabilitation delivered two versus three times per week. *Spine J* 2002;2:402-07.

Rao RD, Smuck M. *In Orthopaedic Knowledge Update: Spine 4*. 2012.

Ray AL. Neuroplasticity, sensitization, and pain. *In Comprehensive Treatment of Chronic Pain by Medical, Interventional, and Integrative Approaches*. Chapter 71. 2012.

Reisfield GM, Goldberger BA, Bertholf RL. 'False-positive' and 'false-negative' test results in clinical urine drug testing. *Bioanalysis* 2009;1:937-52.

Reuben DB, Alvanzo AA, Ashikaga T, et al. National institutes of health pathways to prevention workshop: the role of opioids in the treatment of chronic pain. *Ann Intern Med* 2015;162:295-300.

Richards BL, Whittle SL, van der Heijde DM, Buchbinder R. The efficacy and safety of muscle relaxants in inflammatory arthritis: a Cochrane systematic review. *J Rheumatol Suppl* 2012;90:34-39.

Riew KD, Park JB, Cho YS, et al. Nerve root blocks in the treatment of lumbar radicular pain: a minimum five-year follow-up. *J Bone Joint Surg Am* 2006;88:1722-25.

Robinson MJ, Sheehan D, Gaynor PJ, et al. Relationship between major depressive disorder and associated painful physical symptoms: analysis of data from two pooled placebo-controlled, randomized studies of duloxetine. *Int Clin Psychopharmacol* 2013;28:330-38.

Rolita L, Spegman A, Tang X, Cronstein BN. Increasing narcotic analgesic prescriptions for osteoarthritis is associated with increased falls and fractures in the elderly. *J Am Geriatr Soc* 2013;61:335-40.

Romera I, Pérez V, Menchón JM, et al. Early vs. conventional switching of antidepressants in patients with MDD and moderate to severe pain: a double-blind randomized study. *J Affect Disord* 2012;143:47-55.

Ross JR, Rutter D, Welsh K, et al. Clinical response to morphine in cancer patients and genetic variation in candidate genes. *Pharmacogenomics J* 2005;5:324-36.

Ross SM. Osteoarthritis: a proprietary Arnica gel is found to be as effective as ibuprofen gel in osteoarthritis of the hands. *Holist Nurs Pract* 2008;22:237-39.

Rowbotham MC, Davies PS, Fields HL. Topical lidocaine gel relieves postherpetic neuralgia. *Ann Neurol* 1995;37:246-53.

Rubin R. Analysis reveals large increase in hospitalizations in recent years among older patients prescribed opioids. *JAMA* 2014;312:1621-23.

Rutkow L, Chang HY, Daubresse M, et al. Effect of Florida's prescription drug monitoring program and pill mill laws on opioid prescribing and use. *JAMA Intern Med* 2015;175:1642-49.

Sansone RA, Sansone LA. Chronic pain syndromes and borderline personality. *Innov Clin Neurosci* 2012;9:10-14.

Santilli V, Beghi E, Finucci S. Chiropractic manipulation in the treatment of acute back pain and sciatica with disc protrusion: a randomized double-blind clinical trial of active and simulated spinal manipulations. *Spine J* 2006;6:131-37.

Sarzi-Puttini P, Atzeni F, Lanata L, Bagnasco M. Efficacy of ketoprofen vs. ibuprofen and diclofenac: a systematic review of the literature and meta-analysis. *Clin Exp Rheumatol* 2013;31:731-38.

[Return to Table of Contents](#)

References

- Saunders KW, Dunn KM, Merrill JO, et al. Relationship of opioid use and dosage levels to fractures in older chronic pain patients. *J Gen Intern Med* 2010;25:310-15.
- Scherder E, Oosterman J, Swaab D, et al. Recent developments in pain in dementia. *BMJ* 2005;330:461-64.
- Schisler RE, Groninger H, Rosielle DA. Counseling patients on side effects and driving when starting opioids #248. *J Palliat Med* 2012;15:484-85.
- Scholten PM, Patel SI, Christos PJ, Singh JR. Short-term efficacy of sacroiliac joint corticosteroid injection based on arthrographic contrast patterns. *PM R* 2015;7:385-91.
- Schwartz RP, Gryczynski J, O'Grady KE, et al. Opioid agonist treatments and heroin overdose deaths in Baltimore, Maryland, 1995-2009. *Am J Public Health* 2013;103:917-22.
- Seal KH, Shi Y, Cohen G, et al. Association of mental health disorders with prescription opioids and high-risk opioid use in U.S. veterans of Iraq and Afghanistan. *JAMA* 2012;307:940-47.
- Sehgal N, Manchikanti L, Smith HS. Prescription opioid abuse in chronic pain: a review of opioid abuse predictors and strategies to curb opioid abuse. *Pain Physician* 2012;15:ES67-92.
- Semenchuk MR, Sherman S, Davis B. Double-blind, randomized trial of bupropion SR for the treatment of neuropathic pain. *Neurology* 2001;57:1583-88.
- Sessler NE, Downing JM, Kale H, et al. Reductions in reported deaths following the introduction of extended-release oxycodone (OxyContin) with an abuse-deterrent formulation. *Pharmacoepidemiol Drug Saf* 2014;23:1238-46.
- Severeijns R, Vlaeyen JW, van den Hout MA, Weber WE. Pain catastrophizing predicts pain intensity, disability, and psychological distress independent of the level of physical impairment. *Clin J Pain* 2001;17:165-72.
- Severtson SG, Bartelson BB, Davis JM, et al. 92 difference in rates of abuse following reformulation of extended release oxycodone using data from the RADARS® system poison center program. *Ann Emer Med* 2012;60:S34-35.
- Severtson SG, Bartelson BB, Davis JM, et al. Reduced abuse, therapeutic errors, and diversion following reformulation of extended-release oxycodone in 2010. *J Pain* 2013;14:1122-30.
- Shah A, Hayes CJ, Martin BC. Characteristics of initial prescription episodes and likelihood of long-term opioid use – United States, 2006-2015. *MMWR Morb Mortal Wkly Rep* 2017;66:265-69.
- Shapiro SL, Schwartz GE. Intentional systemic mindfulness: an integrative model for self-regulation and health. *Adv Mind Body Med* 2000;16:128-34.
- Sharp TJ, Harvey AG. Chronic pain and posttraumatic stress disorder: mutual maintenance? *Clin Psychol Rev* 2001;21:857-77.
- Simmons JW, McMillin JN, Emery SF, Kimmich SJ. Intradiscal steroids: a prospective double-blind clinical trial. *Spine* 1992;17:S172-75.
- Simopoulos TT, Manchikanti L, Gupta S, et al. Systematic review of the diagnostic accuracy and therapeutic effectiveness of sacroiliac joint interventions. *Pain Physician* 2015;18:E713-56.
- Simpson EL, Duenas A, Holmes MW, et al. Spinal cord stimulation for chronic pain of neuropathic or ischaemic origin: systematic review and economic evaluation. *Health Technol Assess* 2009;13:iii, ix-x, 1-154.
- Sindrup SH, Bach FW, Madsen C, et al. Venlafaxine versus imipramine in painful polyneuropathy: a randomized, controlled trial. *Neurology* 2003;60:1284-89.

[Return to Table of Contents](#)

References

- Smyth J, Helm R. Focused expressive writing as self-help for stress and trauma. *J Clin Psychol* 2003;59:227-35.
- Snedecor SJ, Sudharshan L, Cappelleri JC, et al. Systematic review and comparison of pharmacologic therapies for neuropathic pain associated with spinal cord injury. *J Pain Res* 2013;6:539-47.
- Solomon DH, Rassen JA, Glynn RJ, et al. The comparative safety of opioids for nonmalignant pain in older adults. *Arch Intern Med* 2010;170:1979-86.
- Spector W, Shaffer T, Potter DE, et al. Risk factors associated with the occurrence of fractures in U.S. nursing homes: resident and facility characteristics and prescription medications. *J Am Geriatr Soc* 2007;55:327-33.
- Spiegel BMR, Younossi ZM, Hays RD, et al. Impact of hepatitis C on health related quality of life: a systematic review and quantitative assessment. *Hepatology* 2005;41:790-800.
- Staal JB, de Bie RA, de Vet HC, et al. Injection therapy for subacute and chronic low back pain: an updated Cochrane review. *Spine* 2009;34:49-59.
- Standaert CJ, Friedly J, Erwin MW, et al. Comparative effectiveness of exercise, acupuncture, and spinal manipulation for low back pain. *Spine* 2011;36:S120-30.
- Stanton T. Supplements for OA: an unconvincing story. *AAOS* 2012.
- Starrels JL, Becker WC, Alford DP, et al. Systematic review: treatment agreements and urine drug testing to reduce opioid misuse in patients with chronic pain. *Ann Intern Med* 2010;152:712-20.
- Starrels JL, Fox AD, Kunins HV, Cunningham CO. They don't know what they don't know: internal medicine residents' knowledge and confidence in urine drug test interpretation for patients with chronic pain. *J Gen Intern Med* 2012;27:1521-27.
- Steenstra IA, Verbeek JH, Heymans MW, Bongers PM. Prognostic factors for duration of sick leave in patients sick listed with acute low back pain: a systematic review of the literature. *Occup Environ Med* 2005;62:851-60.
- Stoffers JM, Völlm BA, Rucker G, et al. Psychological therapies for people with borderline personality disorder. *Cochrane Database Syst Rev* 2012;(8):CD005652.
- Strang J, Manning V, Mayet S, et al. Overdose training and take-home naloxone for opiate users: prospective cohort study of impact on knowledge and attitudes and subsequent management of overdoses. *Addiction* 2008;103:1648-57.
- Substance Abuse and Mental Health Services Administration, The. SAMHSA releases proposed changes to 42 CFR part 2 (alcohol/drug confidentiality rules). Available at: <http://iac.org/samhsa-releases-proposed-changes-to-42-cfr-part-2-alcoholdrug-confidentiality-rules/>. 2016.
- Substance Abuse and Mental Health Services Administration, The. SAMHSA reports highlight rise in tramadol-related hospital emergency department visits. Available at: <http://www.samhsa.gov/newsroom/press-announcements/201505141130>. 2015.
- Sullivan MJ, Thorn B, Haythornthwaite JA, et al. Theoretical perspectives on the relation between catastrophizing and pain. *Clin J Pain* 2001;17:52-64.
- Sun J, Bi J, Chan G, et al. Improved methods to identify stable, highly heritable subtypes of opioid use and related behaviors. *Addict Behav* 2012;37:1138-44.
- Suri P, Saunders KW, Von Korff M. Prevalence and characteristics of flare-ups of chronic non-specific back pain in primary care: a telephone survey. *Clin J Pain* 2012;28:573-80.
- Tait RC, Chibnall JT. Under-treatment of pain in dementia: assessment is key. *J Am Med Dir Assoc* 2008;9:372-74.

References

- Taylor RS, De Vries J, Buchser E, Dejongste MJ. Spinal cord stimulation in the treatment of refractory angina: systematic review and meta-analysis of randomised controlled trials. *BMC Cardiovasc Disord* 2009;9:13.
- Tekur P, Nagarathna R, Chametcha S, et al. A comprehensive yoga programs improves pain, anxiety and depression in chronic low back pain patients more than exercise: an RCT. *Complement Ther Med* 2012;20:107-18.
- Thomas CP, Fullerton CA, Kim M, et al. Medication-assisted treatment with buprenorphine: assessing the evidence. *Psychiatr Serv* 2014;65:158-70.
- Tilbrook HE, Hewitt CE, Aplin JD, et al. Compliance effects in a randomised controlled trial of yoga for chronic low back pain: a methodological study. *Physiotherapy* 2014;100:256-62.
- Tofferi JK, Jackson JL, O'Malley PG. Treatment of fibromyalgia with cyclobenzaprine: a meta-analysis. *Arthritis Rheum* 2004;51:9-13.
- Tolia VN, Patrick SW, Bennett MM, et al. Increasing incidence of the neonatal abstinence syndrome in U.S. neonatal ICUs. *N Engl J Med* 2015;372:2118-26.
- Tramadol (package insert). Amneal Pharmaceuticals, Paterson, NJ; 2012. Available at: <http://druginserts.com/lib/rx/meds/tramadol-hydrochloride-54/>.
- Tramadol ER (package insert). Par Pharmaceutical Companies, Inc., Spring Valley, NY 10977; 2009. Available at: <http://druginserts.com/lib/rx/meds/tramadol-er/>.
- Transmucosal Immediate Release Fentanyl (TIRF) Products Risk Evaluation and Mitigation Strategy (REMS). Education program for prescribers and pharmacists. Accessed July 2016.
- Treatment of painful diabetic neuropathy with topical capsaicin: a multicenter, double-blind, vehicle-controlled study. The Capsaicin Study Group. *Arch Intern Med* 1991;151:2225-29.
- Trescot AM, Datta S, Lee M, Hansen H. Opioid pharmacology. *Pain Physician* 2008;11:S133-53.
- Tsagaridas N, Naka K, Tsapogas P, et al. Spinal cord stimulation in refractory angina: a systematic review of randomized controlled trials. *Acta Cardiol* 2015;70:233-43.
- Turk DC, Gatchel RJ. *In Psychological Approaches to Pain Management, Second Edition: A Practitioner's Handbook*. 2002.
- Turner BJ, Liang Y. Drug overdose in a retrospective cohort with non-cancer pain treated with opioids, antidepressants, and/or sedative-hypnotics: interactions with mental health disorders. *J Gen Intern Med* 2015;30:1081-96.
- Turner JA, Mancl L, Aaron LA. Short- and long-term efficacy of brief cognitive-behavioral therapy for patients with chronic temporomandibular disorder pain: a randomized, controlled trial. *Pain* 2006;121:181-94.
- Turner JA, Sears JM, Loeser JD. Programmable intrathecal opioid delivery systems for chronic noncancer pain: a systematic review of effectiveness and complications. *Clin J Pain* 2007;23:180-95.
- U.S. Department of Health and Human Services. *Ayurvedic medicine*. 2013.
- U.S. Department of Veterans Affairs. *Management of opioid therapy for chronic pain*. 2010.
- U.S. Food and Drug Administration. FDA reminds the public about the potential for life-threatening harm from accidental exposure to fentanyl transdermal systems ("patches"). Available at: <http://www.fda.gov/drugs/drugsafety/ucm300747.htm>. 2012.

[Return to Table of Contents](#)

References

- U.S. Food and Drug Administration. Fentanyl patch can be deadly to children. Available at: <http://www.fda.gov/forconsumers/consumerupdates/ucm300803.htm>. 2013.
- U.S. Food and Drug Administration. How to dispose of unused medications. 2015.
- Ussher M, Spatz A, Copland C, et al. Immediate effects of a brief mindfulness-based body scan on patients with chronic pain. *J Behav Med* 2014;37:127-34.
- Vadivelu N, Huang Y, Mirante B, et al. Patient considerations in the use of tapentadol for moderate to severe pain. *Drug Healthc Patient Saf* 2013;5:151-59.
- van Dam van Isselt EF, Groenewegen-Sipkema KH, Spruit-van Eijk M, et al. Pain in patients with COPD: a systematic review and meta-analysis. *BMJ Open* 2014;4:e005898.
- van Middelkoop M, Rubinstein SM, Kuijpers T, et al. A systematic review on the effectiveness of physical and rehabilitation interventions for chronic non-specific low back pain. *Eur Spine J* 2011;20:19-39.
- van Tulder MW, Touray T, Furlan AD, et al. Muscle relaxants for non-specific low back pain. *Cochrane Database Syst Rev* 2003;(2):CD004252.
- Vargas-Espinosa ML, Sanmartí-García G, Vázquez-Delgado E, Gay-Escoda C. Antiepileptic drugs for the treatment of neuropathic pain: a systematic review. *Med Oral Patol Oral Cir Bucal* 2012;17:e786-93.
- Vestergaard P, Rejnmark L, Mosekilde L. Fracture risk associated with the use of morphine and opiates. *J Intern Med* 2006;260:76-87.
- Vincent K, Maigne JY, Fischhoff C, et al. Systematic review of manual therapies for non-specific neck pain. *Joint Bone Spine* 2013;80:508-15.
- Vissers KC, Besse K, Hans G, et al. Opioid rotation in the management of chronic pain: where is the evidence? *Pain Pract* 2010;10:85-93.
- Viteri OA, Soto EE, Bahado-Singh RO, et al. Fetal anomalies and long-term effects associated with substance abuse in pregnancy: a literature review. *Am J Perinatol* 2015;32:405-16.
- Vitton O, Gendreau M, Gendreau J, et al. A double-blind placebo-controlled trial of milnacipran in the treatment of fibromyalgia. *Hum Psychopharmacol* 2004;19:S27-35.
- Vlaeyen JW, de Jong J, Geilen M, et al. The treatment of fear of movement/(re)injury in chronic low back pain: further evidence on the effectiveness of exposure in vivo. *Clin J Pain* 2002;18:251-61.
- Voelker R. Sharp increase in women's deaths from Rx painkiller overdoses. *JAMA* 2013;310:789.
- Volkow ND. Opioids in pregnancy. *BMJ* 2016a;352:i19.
- Volkow ND, Frieden TR, Hyde PS, Cha SS. Medication-assisted therapies – tackling the opioid-overdose epidemic. *N Engl J Med* 2014;370:2063-66.
- Volkow ND, McLellan AT. Opioid abuse in chronic pain – misconceptions and mitigation strategies. *N Engl J Med* 2016b;374:1253-63.
- Volkow ND, McLellan TA. Curtailing diversion and abuse of opioid analgesics without jeopardizing pain treatment. *JAMA* 2011;305:1346-47.
- Von Korff M, Kolodny A, Deyo RA, Chou R. Long-term opioid therapy reconsidered. *Ann Intern Med* 2011;155:325-28.
- Vowles KE, McEntee ML, Julnes PS, et al. Rates of opioid misuse, abuse, and addiction in chronic pain: a systematic review and data synthesis. *Pain* 2015;156:569-76.

[Return to Table of Contents](#)

References

Walker BF, French SD, Grant W, Green S. Combined chiropractic interventions for low-back pain (review). *Cochrane Database Syst Rev* 2010:CD005427.

Wallace LS, Dhingra LK. A systematic review of smartphone applications for chronic pain available for download in the United States. *J Opioid Manag* 2014;10:63-68.

Wallace MS, Marcotte TD, Umlauf A, et al. Efficacy of inhaled cannabis on painful diabetic neuropathy. *J Pain* 2015;16:616-27.

Wang J, Christo PJ. The influence of prescription monitoring programs on chronic pain management. *Pain Physician* 2009;12:507-15.

Wardell DW, Decker SA, Engebretson JC. Healing touch for older adults with persistent pain. *Holist Nurs Pract* 2012;26:194-202.

Wasan AD, Butler SF, Budman SH, et al. Psychiatric history and psychologic adjustment as risk factors for aberrant drug-related behavior among patients with chronic pain. *Clin J Pain* 2007;23:307-15.

Wasan AD, Michna E, Edwards RR, et al. Psychiatric comorbidity is associated prospectively with diminished opioid analgesia and increased opioid misuse in patients with chronic low back pain. *Anesthesiology* 2015;123:861-72.

Waseem Z, Boulias C, Gordon A, et al. Botulinum toxin injections for low-back pain and sciatica. *Cochrane Database Syst Rev* 2011;(1):CD008257.

Washington State – Agency Medical Directors' Group. Interagency guideline on prescribing opioids for pain. Third Edition. June 2015.

Webster LR, Slevin KA, Narayana A, et al. Fentanyl buccal tablet compared with immediate-release oxycodone for the management of breakthrough pain in opioid-tolerant patients with chronic cancer and noncancer pain: a randomized, double-blind, crossover study followed by a 12-week open-label phase to evaluate patient outcomes. *Pain Med* 2013;14:1332-45.

Webster LR, Webster RM. Predicting aberrant behaviors in opioid-treated patients: preliminary validation of the opioid risk tool. *Pain Med* 2005;6:432-42.

Weiner RS. *In Pain Management: A Practical Guide for Clinicians, Sixth Edition.* Amer Acad Pain Mgmt 2001.

Wesson DR, Ling W. The clinical opiate withdrawal scale (COWS). *J Psychoactive Drugs* 2003;35:253-59.

Whiteman VE, Salemi JL, Mogos MF, et al. Maternal opioid drug use during pregnancy and its impact on perinatal morbidity, mortality, and the costs of medical care in the United States. *J Pregnancy* 2014;2014:906723.

Wiffen PJ, Derry S, Moore RA, et al. Antiepileptic drugs for neuropathic pain and fibromyalgia – an overview of Cochrane reviews. *Cochrane Database Syst Rev* 2013;(11):CD010567.

Wiffen PJ, Rees J. Lamotrigine for acute and chronic pain. *Cochrane Database Syst Rev* 2007;(2):CD006044.

Winslow LC, Kroll DJ. Herbs as medicines. *Arch Intern Med* 1998;158:2192-99.

Wisconsin Medical Society Task Force on Pain Management. Guidelines for the assessment and management of chronic pain. *WMJ* 2004;103:13-42.

Wong E, Walker KA. A review of common methods to convert morphine to methadone. *J Community Hosp Intern Med Perspect* 2013;2.

[Return to Table of Contents](#)