

References

- Aasvang EK, Lunn TH, Hansen TB. Chronic pre-operative opioid use and acute pain after fast-track total knee arthroplasty. *Acta Anaesthesiol Scand*. 2016 Apr;60(4):529-36.
- Alam A, Gomes T, Zheng H, et al. Long-term analgesic use after low-risk surgery: a retrospective cohort study. *Arch Intern Med*. 2012 Mar 12;172(5):425-30.
- Alford DP, Compton P, Samet JH. Acute pain management for patients receiving maintenance methadone or buprenorphine therapy. *Ann Intern Med*. 2006 Jan 17;144(2):127-34.
- American College of Obstetricians and Gynecologists. ACOG Committee Opinion No. 775. Nonobstetric surgery during pregnancy. *Obstet Gynecol* 2019;133:e285–6.
- American Diabetes Association. 15. Diabetes Care in the Hospital: Standards of Medical Care in Diabetes—2019. *Diabetes Care* 2019 Jan; 42(Supplement 1): S173-S181.
- American Society of Anesthesiologists. Pregnancy Testing Prior to Anesthesia and Surgery. October 26, 2016.
- American Society of Anesthesiologists Committee. Practice guidelines for preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration: application to healthy patients undergoing elective procedures: an updated report by the American Society of Anesthesiologists Committee on Standards and Practice Parameters. *Anesthesiology* 2017; 126:376–93.
- Appfelbaum JL, Ashburn MA, Connis RT, et al. Practice Guidelines for Acute Pain Management in the Perioperative Setting: An Updated Report by the American Society of Anesthesiologists Task Force on Acute Pain Management. *Anesthesiology* 2012a; 116:248-73.
- Appfelbaum JL, Connis RT, Nickinovich DG. Practice advisory for preanesthesia evaluation: An updated report by the American Society of Anesthesiologists Task Force on Preanesthesia Evaluation. *Anesthesiology* 2012b; 116:522–38.
- Babu KM, Brent J, Juurlink DN. Prevention of Opioid Overdose. *N Engl J Med*. 2019 Jun 6;380(23):2246-2255.
- Barry DT, Beitel M, Garnet B, et al. Relations among psychopathology, substance use, and physical pain experiences in methadone-maintained patients. *J Clin Psychiatry*. 2009 Sep; 70(9): 1213–1218.
- Bateman BT, Cole NM, Maeda A, et al. Patterns of Opioid Prescription and Use After Cesarean Delivery. *Obstet Gynecol* 2017;130(1):29-35.
- Bates C, Laciak R, Southwick A, Bishoff J. Over-prescription of postoperative narcotics: a look at postoperative pain medication delivery, consumption and disposal in urological practice. *The J Urol* 2011;185(2):551-5.
- Bayfield NGR, Pannekoek A, Tian DH. Preoperative cigarette smoking and short-term morbidity and mortality after cardiac surgery: a meta-analysis. *Heart Asia*. 2018 Oct 24;10(2):e011069.
- Bedard NA, Sierra RJ, Mabry T. Opioids After Orthopaedic Surgery: There Is a Need for Universal Prescribing Recommendations: Commentary on an article by Matthew J. Sabatino, MD, MS, et al.: “Excess Opioid Medication and Variation in Prescribing Patterns Following Common Orthopaedic Procedures”. *J Bone Joint Surg Am*. 2018 Feb 7;100(3):e17.
- Behzadi M, Joukara S, Beikb. Opioids and Cardiac Arrhythmia: A Literature Review *Med Princ Pract* 2018;27:401–414.
- Bell KL, Shohat N, Goswami K, et al. Preoperative Opioids Increase the Risk of Periprosthetic Joint Infection After Total Joint Arthroplasty. *J Arthroplasty*. 2018 Oct;33(10):3246-3251.e1.

References

- Bentzley BS1, Barth KS2, Back SE3, Book SW4. Discontinuation of buprenorphine maintenance therapy: perspectives and outcomes. *J Subst Abuse Treat*. 2015 May;52:48-57.
- Benyamin R, Trescot AM, Datta S, et al. Opioid complications and side effects. *Pain Physician*. 2008 Mar;11(2 Suppl):S105-20.
- Berglund DD, Rosas S, Kurowicki J, et al. Preoperative Opioid Use Among Patients Undergoing Shoulder Arthroplasty Predicts Prolonged Postoperative Opioid Use. *J Am Acad Orthop Surg*. 2018 Aug 1;27(15):e691-e695.
- Bicket MC, Long JJ, Pronovost PJ, et al. Prescription Opioid Analgesics Commonly Unused After Surgery: A Systemic Review. *JAMA Surg*. 2017 Nov 1;152(11):1066-1071..
- Blessberger H, Kammler J, Domanovits H, et al. Perioperative Beta blockers for preventing surgery-related mortality and morbidity. *Cochrane Database of Systematic Reviews* 2018, Issue 3. Art. No.: CD004476.
- Blevins Peratikos M, Weeks HL, Pisansky AJB et al. Effect of Preoperative Opioid Use on Adverse Outcomes, Medical Spending, and Persistent Opioid Use Following Elective Total Joint Arthroplasty in the United States: A Large Retrospective Cohort Study of Administrative Claims Data. *Pain Med*. 2019 May 23.
- Bock M, Johansson T, Fritsch G, et al. The impact of preoperative testing for blood glucose concentration and haemoglobin A1c on mortality, changes in management and complications in noncardiac elective surgery: a systematic review. *Eur J Anaesthesiol*. 2015 Mar;32(3):152-9.
- Bonner BE, Castillo TN, Fitz DW, et al. Preoperative Opioid Use Negatively Affects Patient-reported Outcomes After Primary Total Hip Arthroplasty. *J Am Acad Orthop Surg*. 2019 Feb 28.
- Boostani R, Derakhshan S. Tramadol induced seizure: A 3-year study. *Caspian J Intern Med*. 2012 Summer;3(3):484-7.
- Brat GA, Agniel D, Beam A, et al. Postsurgical prescriptions for opioid naive patients and association with overdose and misuse: retrospective cohort study. *BMJ* 2018;360:j5790.
- Breen CL, Harris SJ, Lintzeris N, et al. Cessation of methadone maintenance treatment using buprenorphine: transfer from methadone to buprenorphine and subsequent buprenorphine reductions. *Drug and Alcohol Dependence*. Volume 71, Issue 1, 20 July 2003, Pages 49-55.
- Brummett CM, Waljee JF, Goesling J, et al. New Persistent Opioid Use After Minor and Major Surgical Procedures in US Adults. *JAMA Surg*. 2017 Jun 21;152(6):e170504.
- Buchleitner AM, Martínez-Alonso M, Hernández M, Solà I, Mauricio D. Perioperative glycaemic control for diabetic patients undergoing surgery. *Cochrane Database Syst Rev*. 2012 Sep 12;(9):CD007315.
- Buckeridge D, Huang A, Hanley J, et al. Risk of injury associated with opioid use in older adults. *J Am Geriatr Soc*. 2010 Sep;58(9):1664-70.
- Carroll I, Barelka P, Wang CK, et al. A pilot cohort study of the determinants of longitudinal opioid use after surgery. *Anesth Analg*. 2012 Sep;115(3):694-702.
- CDC, 2018. Using Naloxone to Reverse Opioid Overdose in the Workplace: Information for Employers and Workers. DHHS (NIOSH) Publication Number 2019-101. October 2018.
- Chan MTV, Wang CY, Seet E, et al. Association of Unrecognized Obstructive Sleep Apnea with Postoperative Cardiovascular Events in Patients Undergoing Major Noncardiac Surgery. *JAMA*. 2019;321(18):1788-1798.
- Cheah JW, Sing DC, McLaughlin D, et al. The perioperative effects of chronic preoperative opioid use on shoulder arthroplasty outcomes. *J Shoulder Elbow Surg*. 2017 Nov;26(11):1908-1914.

References

- Chung F, Memtsoudis SG, Ramachandran SK, et al. Society of Anesthesia and Sleep Medicine Guidelines on Preoperative Screening and Assessment of Adult Patients with Obstructive Sleep Apnea. *Anesth Analg*. 2016 Aug;123(2):452-73.
- Clark NC, Lintzeris N, Muhleisen PJ. Severe opiate withdrawal in a heroin user precipitated by a massive buprenorphine dose. *Med J Aust*. 2002;176:166–7.
- Clegg A, Young JB. Which medications to avoid in people at risk of delirium: a systematic review. *Age Ageing*. 2011 Jan;40(1):23-9.
- Cron DC, Englesbe MJ, Bolton CJ, et al. Preoperative Opioid Use is Independently Associated With Increased Costs and Worse Outcomes After Major Abdominal Surgery. *Ann Surg*. 2017 Apr;265(4):695-701.
- Davis JM, Lin LA, Liu H, Sites BD. Prescription Opioid Use among Adults with Mental Health Disorders in the United States. *J Am Board Fam Med*. 2017 Jul-Aug;30(4):407-417.
- Dean L. Codeine Therapy and CYP2D6 Genotype. 2012 Sep 20 [Updated 2017 Mar 16]. In: Pratt V, McLeod H, Rubinstein W, et al., editors. Medical Genetics Summaries [Internet]. Bethesda (MD): National Center for Biotechnology Information (US); 2012-. Bookshelf URL: <https://www.ncbi.nlm.nih.gov/books/>.
- Dean, M. Opioids in renal failure and dialysis patients. *J Pain Symptom Manage*. 2004 Nov;28(5):497-504.
- Demma LJ, Carlson KT, Duggan EW, et al. Effect of basal insulin dosage on blood glucose concentration in ambulatory surgery patients with type 2 diabetes. *J Clin Anesth*. 2017 Feb;36:184-188.
- Dhalla IA, Mamdani MM, Siviolotti MLA, et al. Prescribing of opioid analgesics and related mortality before and after the introduction of long-acting oxycodone. *CMAJ*. 2009 Dec 8; 181(12): 891–896.
- Doleman B, Read D, Lund JN, Williams JP. Preventive Acetaminophen Reduces Postoperative Opioid Consumption, Vomiting, and Pain Scores After Surgery: Systematic Review and Meta-Analysis. *Reg Anesth Pain Med*. 2015 Nov-Dec;40(6):706-12.
- Duggan EW, Carlson K, Umpierrez GE. Perioperative Hyperglycemia Management: An Update. *Anesthesiology*. 2017 Mar; 126(3): 547–560.
- Faour M, Anderson JT, Haas AR, et al. Prolonged Preoperative Opioid Therapy Associated With Poor Return to Work Rates After Single-Level Cervical Fusion for Radiculopathy for Patients Receiving Workers' Compensation Benefits. *Spine (Phila Pa 1976)*. 2017 Jan 15;42(2):E104-E110.
- Feinberg AE, Chesney TR, Srikandarajah S. Opioid Use After Discharge in Postoperative Patients: A Systematic Review. *Annals of Surgery*: June 2018 - Volume 267 - Issue 6 - p 1056–1062.
- Fleisher LA, Fleischmann KE, Auerbach AD, et al. 2014 ACC/AHA Guideline on Perioperative Cardiovascular Evaluation and Management of Patients Undergoing Noncardiac Surgery: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. December 2014: 64(22).
- Food and Drug Administration (FDA). FDA Drug Safety Communication: FDA restricts use of prescription codeine pain and cough medicines and tramadol pain medicines in children; recommends against use in breastfeeding women. <https://www.fda.gov/drugs/drug-safety-and-availability/fda-drug-safety-communication-fda-restricts-use-prescription-codeine-pain-and-cough-medicines-and>. Content current as of: March 8, 2018. Accessed: October 20, 2019.
- Food and Drug Administration (FDA). FDA Drug Safety Communication: FDA warns about several safety issues with opioid pain medicines; requires label changes. <https://www.fda.gov/drugs/drug-safety-and-availability/fda-drug-safety-communication-fda-warns-about-several-safety-issues-opioid-pain-medicines-requires?source=govdelivery>. Content current as of: March 30, 2016. Accessed: October 20, 2019.

References

- Fowler AJ, Ahmad T, Phull MK, et al. Meta-analysis of the association between preoperative anaemia and mortality after surgery. *Br J Surg*. 2015 Oct;102(11):1314-24.
- Franco LM, Cota GF, Pinto TS, Ercole FF. Preoperative bathing of the surgical site with chlorhexidine for infection prevention: Systematic review with meta-analysis. *Am J Infect Control*. 2017 Apr 1;45(4):343-349.
- Gandhi GY, Nuttall GA, Abel MD, et al. Intensive intraoperative insulin therapy versus conventional glucose management during cardiac surgery: a randomized trial. *Ann Intern Med*. 2007 Feb 20;146(4):233-43.
- Ganesh A, Maxwell LG. Pathophysiology and management of opioid-induced pruritus. *Drugs*. 2007;67(16):2323-33.
- Gatchel RJ, McGeary DD, McGeary CA, Lippe B. Interdisciplinary chronic pain management: past, present, and future. *Am Psychol*. 2014 Feb-Mar;69(2):119-30.
- Gatchel RJ, Peng YB, Peters ML, et al. The biopsychosocial approach to chronic pain: scientific advances and future directions. *Psychol Bull*. 2007 Jul;133(4):581-624.
- Gatchel RJ, Okifuji A. Evidence-based scientific data documenting the treatment and cost-effectiveness of comprehensive pain programs for chronic nonmalignant pain. *J Pain* 2006;7:779-93.
- Goplen CM, Verbeek W, Kang SH, et al. Preoperative opioid use is associated with worse patient outcomes after Total joint arthroplasty: a systematic review and meta-analysis. *BMC Musculoskeletal Disord*. 2019 May 18;20(1):234.
- Greenwald MK, Comer SD, Fiellin DA. Buprenorphine maintenance and mu-opioid receptor availability in the treatment of opioid use disorder: implications for clinical use and policy. *Drug Alcohol Depend*. 2014 Nov 1;144:1-11.
- Greenwald M, Johanson CE, Bueller J, et al. Buprenorphine duration of action: mu-opioid receptor availability and pharmacokinetic and behavioral indices. *Biol Psychiatry*. 2007 Jan 1;61(1):101-10.
- Gross, JB, Apfelbaum, JI, Caplan, RA, et al. Practice Guidelines for the Perioperative Management of Patients with Obstructive Sleep Apnea: An Updated Report by the American Society of Anesthesiologists Task Force on Perioperative Management of Patients with Obstructive Sleep Apnea. *Anesthesiology*. 2014 Feb;120:268-286.
- Gugelmann HM, Perrone J. Can prescription drug monitoring programs help limit opioid abuse? *JAMA*. 2011 Nov 23;306(20):2258-9.
- Gustafsson UO, Thorell A, Soop M, et al. Haemoglobin A1c as a predictor of postoperative hyperglycaemia and complications after major colorectal surgery. *Br J Surg*. 2009 Nov;96(11):1358-64.
- Hall V, Lynfield R, Hiber WL, et al. Deaths Associated with Opioid Use and Possible Infectious Disease Etiologies Among Persons in the Unexplained Death (UNEX) Surveillance System — Minnesota, 2006–2015. CDC 2018.
- Hansen LE, Stone GL, Matson CA, et al. Total Joint Arthroplasty in Patients Taking Methadone or Buprenorphine/Naloxone Preoperatively for Prior Heroin Addiction: A Prospective Matched Cohort Study. *J Arthroplasty*. 2016 Aug;31(8):1698-701.
- Harrison TK, Kornfeld H, Aggarwal AK, Lembke A. Perioperative Considerations for the Patient with Opioid Use Disorder on Buprenorphine, Methadone, or Naltrexone Maintenance Therapy. *Anesthesiol Clin*. 2018 Sep;36(3):345-359.
- Han B, Compton WM, Blanco C, Jones CM. Prescription Opioid Use, Misuse, and Use Disorders in U.S. Adults. *Ann Intern Med*. 2018 Mar 6;168(5):383-384.

[Return to Table of Contents](#)

References

- Hasin DS, O'Brien C, Auriacombe M, Borges G. DSM-5 Criteria for Substance Use Disorders: Recommendations and Rationale. *American Journal of Psychiatry* 170(8):834-51 August 2013.
- Hassamal S, Haglund M, Wittnebel K, Danovitch I. A preoperative interdisciplinary biopsychosocial opioid reduction program in patients on chronic opioid analgesia prior to spine surgery: A preliminary report and case series. *Scand J Pain*. 2016 Oct;13:27-31.
- Hill MV, Stucke RS, McMahon ML, et al. An Educational Intervention Decreases Opioid Prescribing After General Surgical Operations. *Ann Surg*. 2018 Mar;267(3):468-472.
- Hollmann C, Fernandes NL, Biccadd BM. A Systematic Review of Outcomes Associated with Withholding or Continuing Angiotensin-Converting Enzyme Inhibitors and Angiotensin Receptor Blockers Before Noncardiac Surgery. *Anesth Analg*. 2018 Sep;127(3):678-687.
- Hooten M, Thorson D, Bianco J, Bonte B, Clavel Jr A, Hora J, Johnson C, Kirksson E, Noonan MP, Reznikoff C, Schweim K, Wainio J, Walker N. Institute for Clinical Systems Improvement. Pain: Assessment, Non-Opioid Treatment Approaches and Opioid Management. Updated August 2017.
- Hozack BA, Rivlin M, Lutsky KF, et al. Preoperative Exposure to Benzodiazepines or Sedative/hypnotics Increases the Risk of Greater Filled Opioid Prescriptions After Surgery. *Clin Orthop Relat Res*. 2019 Jun;477(6):1482-1488.
- Issa Y, Ahmed Ali U, Bouwense SA, et al. Preoperative opioid use and the outcome of thoracoscopic splanchnicectomy in chronic pancreatitis: a systematic review. *Surg Endosc*. 2014 Feb;28(2):405-12.
- Jain N, Brock JL, Malik AT, et al. Prediction of Complications, Readmission, and Revision Surgery Based on Duration of Preoperative Opioid Use: Analysis of Major Joint Replacement and Lumbar Fusion. *J Bone Joint Surg Am*. 2019 Mar 6;101(5):384-391.
- Jain N, Brock JL, Phillips FM, et al. Chronic preoperative opioid use is a risk factor for increased complications, resource use, and costs after cervical fusion. *Spine J*. 2018a Nov;18(11):1989-1998.
- Jain N, Phillips FM, Weaver T, Khan SN. Preoperative Chronic Opioid Therapy: A Risk Factor for Complications, Readmission, Continued Opioid Use and Increased Costs After One- and Two-Level Posterior Lumbar Fusion. *Spine (Phila Pa 1976)*. 2018b Oct 1;43(19):1331-1338.
- Jamison RN, Kauffman J, Katz NP. Characteristics of methadone maintenance patients with chronic pain. *J Pain Symptom Manage*. 2000 Jan;19(1):53-62.
- Johansson T, Fritsch G, Flamm M, et al. Effectiveness of non-cardiac preoperative testing in non-cardiac elective surgery: a systematic review. *Br J Anaesth*. 2013 Jun;110(6):926-39.
- Jones HE, O'Grady K, Dahne J, et al. Management of acute postpartum pain in patients maintained on methadone or buprenorphine during pregnancy. *Am J Drug Alcohol Abuse*. 2009;35(3):151-6.
- Joo HS, Wong J, Naik VN, Savoldelli GL. The value of screening preoperative chest X-rays: a systematic review. *Can J Anaesth*. 2005 Jun-Jul;52(6):568-74.
- Kalakoti P, Volkmar AJ, Bedard NA, et al. Preoperative Chronic Opioid Therapy Negatively Impacts Long-term Outcomes Following Cervical Fusion Surgery. *Spine (Phila Pa 1976)*. 2019 Sep;44(18):1279-1286.
- Keay L, Lindsley K, Tielsch J, et al. Routine preoperative medical testing for cataract surgery. *Cochrane Database Syst Rev*. 2019 Jan 8;1:CD007293.
- Kim K, Chen K, Anoushiravani AA, et al. Preoperative Chronic Opioid Use and Its Effects on Total Knee Arthroplasty Outcomes. *J Knee Surg*. 2019 Feb 11.
- Kim KY, Anoushiravani AA, Chen KK, et al. Preoperative Chronic Opioid Users in Total Knee Arthroplasty-Which Patients Persistently Abuse Opiates Following Surgery? *J Arthroplasty*. 2018 Jan;33(1):107-112.

[Return to Table of Contents](#)

References

- Klimas J, Gorfinkel L, Fairbairn N, et al. Strategies to Identify Patient Risks of Prescription Opioid Addiction When Initiating Opioids for Pain: A Systematic Review. *JAMA Netw Open*. 2019 May 3;2(5):e193365.
- Koehler RM, Okoroafor UC, Cannada LK. A systematic review of opioid use after extremity trauma in orthopedic surgery. *Injury*. 2018 Jun;49(6):1003-1007.
- Kork F, Balzer F, Spies CD, et al. Minor Postoperative Increases of Creatinine Are Associated with Higher Mortality and Longer Hospital Length of Stay in Surgical Patients. *Anesthesiology*. 2015 Dec;123(6):1301-1311.
- Kornfeld H, Manfredi L. Effectiveness of full agonist opioids in patients stabilized on buprenorphine undergoing major surgery: a case series. *Am J Ther*. 2010 Sep-Oct;17(5):523-8.
- Kunøe N, Lobmaier P, Vederhus JK, et al. Challenges to antagonist blockade during sustained-release naltrexone treatment. *Addiction*. 2010 Sep;105(9):1633-9.
- Kwon S, Thompson R, Dellinger P, et al. Importance of perioperative glycemic control in general surgery: a report from the Surgical Care and Outcomes Assessment Program. *Ann Surg*. 2013 Jan;257(1):8-14.
- Lambert E, Carey S. Practice Guideline Recommendations on Perioperative Fasting: A Systematic Review. *JPEN J Parenter Enteral Nutr*. 2016 Nov;40(8):1158-1165.
- Larochelle MR, Liebschutz JM, Zhang F, et al. Opioid Prescribing After Nonfatal Overdose and Association with Repeated Overdose: A Cohort Study. *Ann Intern Med*. 2016 Jan 5;164(1):1-9.
- Lee D, Armaghani S, Archer KR1, et al. Preoperative Opioid Use as a Predictor of Adverse Postoperative Self-Reported Outcomes in Patients Undergoing Spine Surgery. *J Bone Joint Surg Am*. 2014 Jun 4;96(11):e89. Epub 2014 Jun 4.
- Lefebvre A, Saliou P, Lucet JC, et al; French Study Group for the Preoperative Prevention of Surgical Site Infections. Preoperative hair removal and surgical site infections: network meta-analysis of randomized controlled trials. *J Hosp Infect*. 2015 Oct;91(2):100-8.
- Lembke A, Ottestad E, Schmiesing C. Patients Maintained on Buprenorphine for Opioid Use Disorder Should Continue Buprenorphine Through the Perioperative Period. *Pain Med*. 2019 Mar 1;20(3):425-428.
- Leung JM, Sands LP, Paul S, et al. Does postoperative delirium limit the use of patient-controlled analgesia in older surgical patients? *Anesthesiology*. 2009 Sep;111(3):625-31.
- Levine GN, Bates ER, Bittl JA, et al. 2016 ACC/AHA Guideline Focused Update on Duration of Dual Antiplatelet Therapy in Patients With Coronary Artery Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines: An Update of the 2011 ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention, 2011 ACCF/AHA Guideline for Coronary Artery Bypass Graft Surgery, 2012 ACC/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the Diagnosis and Management of Patients With Stable Ischemic Heart Disease, 2013 ACCF/AHA Guideline for the Management of ST-Elevation Myocardial Infarction, 2014 AHA/ACC Guideline for the Management of Patients With Non-ST-Elevation Acute Coronary Syndromes, and 2014 ACC/AHA Guideline on Perioperative Cardiovascular Evaluation and Management of Patients Undergoing Noncardiac Surgery. *Circulation*. 2016 Sep 6;134(10):e123-55.
- Levine M. In older adults with COPD, new opioid use was linked to increased risk for respiratory and all-cause mortality. *Ann Intern Med*. 2017 Jan 17;166(2):JC11.
- Ling Q, Gu Y, Chen J, et al. Consequences of continuing renin angiotensin aldosterone system antagonists in the preoperative period: a systematic review and meta-analysis. *BMC Anesthesiol*. 2018 Feb 26;18(1):26.
- Ling W, Hillhouse M, Domier C, et al. Buprenorphine tapering schedule and illicit opioid use. *Addiction*. 2009 Feb;104(2):256-65.

References

- Macintyre PE, Russell RA, Usher KA, et al. Pain relief and opioid requirements in the first 24 hours after surgery in patients taking buprenorphine and methadone opioid substitution therapy. *Anaesth Intensive Care*. 2013 Mar;41(2):222-30.
- Mador MJ, Henderson J. Effect of Opioids on Sleep and Breathing in Chronic Pain Patients. *J Clin Sleep Med*. 2014 Aug 15; 10(8): 853–854.
- Maher JL, Mahabir RC. Preoperative pregnancy testing. *Can J Plast Surg*. 2012 Autumn; 20(3): e32–e34.
- Manninen PH. Opioids and seizures. *Can J Anaesth*. 1997 May;44(5 Pt 1):463-6.
- Mark J, Argentieri DM, Gutierrez CA, et al. Ultrarestrictive Opioid Prescription Protocol for Pain Management After Gynecologic and Abdominal Surgery. *JAMA Netw Open*. 2018 Dec 7;1(8):e185452.
- McNeely J, Wu LT, Subramaniam G, et al. Performance of the Tobacco, Alcohol, Prescription Medication, and Other Substance Use (TAPS) Tool for Substance Use Screening in Primary Care Patients. *Ann Intern Med*. 2016 Nov 15;165(10):690-699.
- Mehta V, Langford RM. Acute pain management for opioid dependent patients. *Anaesthesia*. 2006 Mar;61(3):269-76.
- Meyer M, Paranya G, Keefer Norris A, Howard D. Intrapartum and postpartum analgesia for women maintained on buprenorphine during pregnancy. *Eur J Pain*. 2010 Oct;14(9):939-43.
- Miech R, Johnston L, O'Malley PM, et al. Prescription Opioids in Adolescence and Future Opioid Misuse. *Pediatrics*. 2015 Nov;136(5):e1169-77.
- Miller M, Barber CW, Leatherman S, et al. Prescription opioid duration of action and the risk of unintentional overdose among patients receiving opioid therapy. *JAMA Intern Med*. 2015 Apr;175(4):608-15.
- Miller M, Sturmer T, Azrael D, et al. Opioid analgesics and the risk of fractures in older adults with arthritis. *J Am Geriatr Soc*. 2011 Mar;59(3):430-8.
- Morrison RS, Magaziner J, Gilbert M, et al. Relationship between pain and opioid analgesics on the development of delirium following hip fracture. *J Gerontol A Biol Sci Med Sci*. 2003 Jan;58(1):76-81.
- Munro J, Booth A, Nicholl J. Routine preoperative testing: a systematic review of the evidence. *Health Technol Assess*. 1997;1(12):i-iv; 1-62.
- Nair BG, Horibe M, Neradilek MB, et al. The Effect of Intraoperative Blood Glucose Management on Postoperative Blood Glucose Levels in Noncardiac Surgery Patients. *Anesth Analg*. 2016 Mar;122(3):893-902.
- National Institute for Health and Care Excellence (NICE): Clinical Guidelines. Preoperative Tests (Update): Routine Preoperative Tests for Elective Surgery. London: National Institute for Health and Care Excellence (UK); April 2016.
- Nguyen LC, Sing DC, Bozic KJ. Preoperative Reduction of Opioid Use Before Total Joint Arthroplasty. *J Arthroplasty*. 2016 Sep;31(9 Suppl):282-7.
- Nishimura RA, Otto CM, Bonow RO, et al. 2017 AHA/ACC Focused Update of the 2014 AHA/ACC Guideline for the Management of Patients with Valvular Heart Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2017 Jun 20;135(25):e1159-e1195.
- Nolan MB, Martin DP, Thompson R, et al. Association Between Smoking Status, Preoperative Exhaled Carbon Monoxide Levels, and Postoperative Surgical Site Infection in Patients Undergoing Elective Surgery. *JAMA Surg*. 2017 May 1;152(5):476-483.
- Nolan MB, Warner DO. Safety and Efficacy of Nicotine Replacement Therapy in the Perioperative Period: A Narrative Review. *Mayo Clin Proc*. 2015 Nov;90(11):1553-61.

References

- Olfson M, Crystal S, Wall M, et al. Causes of Death After Nonfatal Opioid Overdose. *JAMA Psychiatry*. 2018 Aug 1;75(8):820-827.
- Oslund S, Robinson RC, Clark TC. Long-term effectiveness of a comprehensive pain management program: strengthening the case for interdisciplinary care. *Proc (Bayl Univ Med Cent)*. 2009 Jul; 22(3): 211–214.
- Osmundson, SS, Schornack LA, Grasc JL, et.al. Post-discharge opioid use after cesarean delivery. *Obstet Gynecol* 2017;130(1)36-41.
- Pedersen PU, Larsen P, Håkonsen SJ. The effectiveness of systematic perioperative oral hygiene in reduction of postoperative respiratory tract infections after elective thoracic surgery in adults: a systematic review. *JBI Database System Rev Implement Rep*. 2016 Jan;14(1):140-73.
- Pham PC, Khaing K, Sievers TM. 2017 update on pain management in patients with chronic kidney disease. *Clin Kidney J*. 2017 Oct; 10(5): 688–697.
- Pierce M, Bird SM, Hickman M, et al. Impact of treatment for opioid dependence on fatal drug-related poisoning: a national cohort study in England. *Addiction*. 2016 Feb;111(2):298-308.
- Politzer CS, Kildow BJ, Goltz DE, et al. Trends in Opioid Utilization Before and After Total Knee Arthroplasty. *J Arthroplasty*. 2018 Jul;33(7S):S147-S153.e1.
- Powell MZ, Mueller SW, Reynolds PM. Assessment of Opioid Cross-reactivity and Provider Perceptions in Hospitalized Patients With Reported Opioid Allergies. *Ann Pharmacother*. 2019 Nov;53(11):1117-1123.
- Quaye AN, Zhang Y. Perioperative Management of Buprenorphine: Solving the Conundrum. *Pain Med*. 2018 Nov 30.
- Quinlan J, Cox F. Acute pain management in patients with drug dependence syndrome. *Pain Rep*. 2017 Jul; 2(4): e611.
- Ramos M, Khalpey Z, Lipsitz S, et al. Relationship of perioperative hyperglycemia and postoperative infections in patients who undergo general and vascular surgery. *Ann Surg*. 2008 Oct;248(4):585-91.
- Ray WA, Chung CP, Murray KT, et al. Prescription of Long-Acting Opioids and Mortality in Patients With Chronic Noncancer Pain. *JAMA*. 2016 Jun 14;315(22):2415-23.
- Rozell JC, Courtney PM, Dattilo JR, et al. Preoperative Opiate Use Independently Predicts Narcotic Consumption and Complications After Total Joint Arthroplasty. *J Arthroplasty*. 2017 Sep;32(9):2658-2662.
- SAMHSA, 2019. <https://www.samhsa.gov/medication-assisted-treatment/treatment/buprenorphine>.
- SAMHSA, 2018. <https://www.store.samhsa.gov/product/Opioid-Overdose-Prevention-Toolkit/SMA18-4742>.
- SAMHSA, 2009. Treatment Improvement Protocol (TIP) Series, No. 49. Report No.: (SMA) 09-4380. <https://www.ncbi.nlm.nih.gov/books/NBK64041>.
- Sathya B, Davis R, Taveira T, et al. Intensity of peri-operative glycemic control and postoperative outcomes in patients with diabetes: a meta-analysis. *Diabetes Res Clin Pract*. 2013 Oct;102(1):8-15.
- Schwarze ML, Barnato AE, Rathouz PJ, et al. Development of a list of high-risk operations for patients 65 years and older. *JAMA Surg*. 2015 Apr;150(4):325-31.
- Seamans MJ, Carey TS, Westreich DJ, et al. Association of Household Opioid Availability and Prescription Opioid Initiation Among Household Members. *JAMA Intern Med*. 2018 Jan 1;178(1):102-109.
- Sen S1, Arulkumar S1, Cornett EM1, et al. New Pain Management Options for the Surgical Patient on Methadone and Buprenorphine. *Curr Pain Headache Rep*. 2016 Mar;20(3):16.

[Return to Table of Contents](#)

References

- Shafi S, Collinsworth AW, Copeland LA, et al. Association of Opioid-Related Adverse Drug Events with Clinical and Cost Outcomes Among Surgical Patients in a Large Integrated Health Care Delivery System. *JAMA Surg*. 2018 Aug 1;153(8):757-763.
- Shah A, Hayes CJ, Martin BC, et al. Characteristics of Initial Prescription Episodes and Likelihood of Long-Term Opioid Use — United States, 2006–2015. *MMWR Weekly* / March 17, 2017 / 66(10);265–269.
- Shi D, Yao Y, Yu W. Comparison of preoperative hair removal methods for the reduction of surgical site infections: a meta-analysis. *J Clin Nurs*. 2017 Oct;26(19-20):2907-2914.
- Sigmon SC1, Dunn KE, Saulsgiver K, et al. A randomized, double-blind evaluation of buprenorphine taper duration in primary prescription opioid abusers. *JAMA Psychiatry*. 2013 Dec;70(12):1347-54.
- Simha V, Shah P. Perioperative Glucose Control in Patients with Diabetes Undergoing Elective Surgery. *JAMA*. 2019;321(4):399-400.
- Solomon DH, Rassen JA, Glynn RJ, et al. The comparative safety of analgesics in older adults with arthritis. *Arch Intern Med*. 2010 Dec 13;170(22):1968-76.
- Strain EC, Preston KL, Liebson IA, Bigelow GE. Buprenorphine effects in methadone-maintained volunteers: effects at two hours after methadone. *J Pharmacol Exp Ther*. 1995;272:628–38.
- Strain EC, Preston KL, Liebson IA, Bigelow GE. Precipitated withdrawal by pentazocine in methadone-maintained volunteers. *J Pharmacol Exp Ther*. 1993;267:624–34.
- Sun EC, Damall BD, Baker LC, Mackey S. Incidence of and Risk Factors for Chronic Opioid Use Among Opioid-Naive Patients in the Postoperative Period. *JAMA Intern Med*. 2016 Sep 1;176(9):1286-93.
- Swart LM, van der Zanden V, Spies PE, et al. The Comparative Risk of Delirium with Different Opioids: A Systematic Review. *Drugs Aging*. 2017 Jun;34(6):437-443.
- Swegle JM, Logemann C. Management of common opioid-induced adverse effects. *Am Fam Physician*. 2006 Oct 15;74(8):1347-54.
- Tan WH, Yu J, Feaman S, et al. Opioid Medication Use in the Surgical Patient: An Assessment of Prescribing Patterns and Use. August 2018 Volume 227, Issue 2, Pages 203–211.
- Thiels CA, Habermann EB, Hooten WM, Jeffery MM. Chronic use of tramadol after acute pain episode: cohort study. *BMJ* 2019;365:l1849.
- Thiels CA, Anderson SS, Ubl DS, et al. Wide Variation and Overprescription of Opioids After Elective Surgery. *Ann Surg* 2017 Oct;266(4):564-573.
- Thomsen T, Villebro N, Møller AM. Interventions for preoperative smoking cessation. *Cochrane Database Syst Rev*. 2014 Mar 27;(3):CD002294.
- Ti L, Voon P, Dobrer S, et al. Denial of pain medication by health care providers predicts in-hospital illicit drug use among individuals who use illicit drugs. *Pain Res Manag*. 2015 Mar-Apr;20(2):84-8. Epub 2015 Jan 6.
- Tisdale JE. Drug-induced QT interval prolongation and torsades de pointes: Role of the pharmacist in risk assessment, prevention and management. *Can Pharm J (Ott)*. 2016 May; 149(3): 139–152.
- Tissot TA. Probable Meperidine-Induced Serotonin Syndrome in a Patient with a History of Fluoxetine Use. *Anesthesiology* 6 2003, Vol.98, 1511-1512.
- Tye EY, Anderson JT, Faour M, et al. Prolonged Preoperative Opioid Therapy in Patients With Degenerative Lumbar Stenosis in a Workers' Compensation Setting. *Spine (Phila Pa 1976)*. 2017 Oct 1;42(19):E1140-E1146.

[Return to Table of Contents](#)

References

- Umpierrez, GE, Smiley, D, Jacobs, S et al. Randomized Study of Basal-Bolus Insulin Therapy in the Inpatient Management of Patients with Type 2 Diabetes Undergoing General Surgery (RABBIT 2 Surgery). *Diabetes Care* 2011 Feb; 34(2): 256-261.
- U.S. Preventive Services Task Force (USPSTF). Abnormal Blood Glucose and Type 2 Diabetes Mellitus: Screening. Release Date: October 2015.
- van den Boom W, Schroeder RA, Manning MW, et al. Effect of A1C and Glucose on Postoperative Mortality in Noncardiac and Cardiac Surgeries. *Diabetes Care*. 2018 Apr;41(4):782-788.
- Vickers AP. Naltrexone and problems in pain management: How to manage acute pain in people taking an opioid antagonist. *BMJ*. 2006 Jan 21; 332(7534): 132–133.
- Vilkins AL, Bagley SM, Hahn KA, et al. Comparison of Post-Cesarean Section Opioid Analgesic Requirements in Women With Opioid Use Disorder Treated With Methadone or Buprenorphine. *J Addict Med*. 2017 Sep/Oct;11(5):397-401.
- Volkow ND, McLellan AT. Opioid Abuse in Chronic Pain--Misconceptions and Mitigation Strategies. *N Engl J Med*. 2016 Mar 31;374(13):1253-63.
- Volkow ND, Frieden TR, Hyde PS, Cha SS. Medication-assisted therapies--tackling the opioid-overdose epidemic. *N Engl J Med*. 2014 May 29;370(22):2063-6.
- Volpe DA, McMahon Tobin GA, Mellon RD, et al. Uniform assessment and ranking of opioid μ receptor binding constants for selected opioid drugs. *Regul Toxicol Pharmacol*. 2011 Apr;59(3):385-90.
- Voon P, Greer AM, Amlani A, et al. Pain as a risk factor for substance use: a qualitative study of people who use drugs in British Columbia, Canada. *Harm Reduct J*. 2018; 15: 35.
- Vozoris NT, Wang X, Fischer HD, et al. Incident opioid drug use and adverse respiratory outcomes among older adults with COPD. *Eur Respir J*. 2016 Sep;48(3):683-93.
- Wahr JA, Parks R, Boisvert D, et al. Preoperative serum potassium levels and perioperative outcomes in cardiac surgery patients. Multicenter Study of Perioperative Ischemia Research Group. *JAMA*. 1999 Jun 16;281(23):2203-10.
- Waljee JF, Cron DC, Steiger RM, et al. Effect of Preoperative Opioid Exposure on Healthcare Utilization and Expenditures Following Elective Abdominal Surgery. *Ann Surg*. 2017 Apr;265(4):715-721.
- Walsh SL, Eissenberg T. The clinical pharmacology of buprenorphine: extrapolating from the laboratory to the clinic. *Drug Alcohol Depend*. 2003;70:S13–27.
- Walsh SL, June HL, Schuh KJ, Preston KL, Bigelow GE, Stitzer ML. Effects of buprenorphine and methadone in methadone-maintained subjects. *Psychopharmacology (Berl)* 1995;119:268–76.
- Webster J, Osborne S. Preoperative bathing or showering with skin antiseptics to prevent surgical site infection. *Cochrane Database Syst Rev*. 2015 Feb 20;(2):CD004985.
- Wiese AD, Griffin MR, Schaffner W, et al. Long-acting Opioid Use and the Risk of Serious Infections: A Retrospective Cohort Study. *Clin Infect Dis*. 2019 May 17;68(11):1862-1869.
- Wiese AD, Griffin MR, Schaffner W, et al. Opioid Analgesic Use and Risk for Invasive Pneumococcal Diseases: A Nested Case-Control Study. *Ann Intern Med*. 2018 Mar 20;168(6):396-404.
- Wijesundera DN, Beattie WS. Calcium channel blockers for reducing cardiac morbidity after noncardiac surgery: a meta-analysis. *Anesth Analg*. 2003 Sep;97(3):634-41.

[Return to Table of Contents](#)

References

Wilson W, Taubert KA, Gewitz M, et al; American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee; American Heart Association Council on Cardiovascular Disease in the Young; American Heart Association Council on Clinical Cardiology; American Heart Association Council on Cardiovascular Surgery and Anesthesia; Quality of Care and Outcomes Research Interdisciplinary Working Group. Prevention of infective endocarditis: guidelines from the American Heart Association: a guideline from the American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee, Council on Cardiovascular Disease in the Young, and the Council on Clinical Cardiology, Council on Cardiovascular Surgery and Anesthesia, and the Quality of Care and Outcomes Research Interdisciplinary Working Group. *Circulation*. 2007 Oct 9;116(15):1736-54.

Yang MH, Jaeger M, Baxter M, et al. Postoperative dysglycemia in elective non-diabetic surgical patients: a prospective observational study. *Can J Anaesth*. 2016 Dec;63(12):1319-1334.

Zarling BJ, Sikora-Klak J, Bergum C, Markel DC. How Do Preoperative Medications Influence Outcomes After Total Joint Arthroplasty? *J Arthroplasty*. 2017 Sep;32(9S):S259-S262.

Zarling BJ, Yokhana SS, Herzog DT, Markel DC. Preoperative and Postoperative Opiate Use by the Arthroplasty Patient. *J Arthroplasty*. 2016 Oct;31(10):2081-4.

Zedler BK, Saunders WB, Joyce AR, et al. Validation of a Screening Risk Index for Serious Prescription Opioid-Induced Respiratory Depression or Overdose in a US Commercial Health Plan Claims Database. *Pain Med*. 2018 Jan 1;19(1):68-78.

[Return to Table of Contents](#)